

Product datasheet for RC212859L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Keratin 80 (KRT80) (NM_182507) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Keratin 80 (KRT80) (NM 182507) Human Tagged ORF Clone Lentiviral Particle

Symbol: Keratin 80

Synonyms: KB20

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

ACCN: NM_182507 **ORF Size:** 1356 bp

ORF Nucleotide

- - - - |

144501

Sequence:
OTI Disclaimer:

Locus ID:

The ORF insert of this clone is exactly the same as(RC212859).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 182507.1</u>

RefSeq Size: 3411 bp

RefSeq ORF: 1359 bp

UniProt ID: Q6KB66

Cytogenetics: 12q13.13

MW: 50.3 kDa





Gene Summary:

Keratins are intermediate filament proteins responsible for the structural integrity of epithelial cells and are subdivided into epithelial keratins and hair keratins. This gene's expression profile shows that it encodes a type II epithelial keratin, although structurally the encoded protein is more like a type II hair keratin. This protein is involved in cell differentiation, localizing near desmosomal plaques in earlier stages of differentiation but then dispersing throughout the cytoplasm in terminally differentiating cells. The type II keratins are clustered in a region of chromosome 12q13. Two transcript variants encoding two different fully functional isoforms have been found for this gene.[provided by RefSeq, Oct 2010]