

## Product datasheet for RC212892L4V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **CPNE1 (NM\_152925) Human Tagged ORF Clone Lentiviral Particle**

**Product data:** 

Product Type: Lentiviral Particles

**Product Name:** CPNE1 (NM\_152925) Human Tagged ORF Clone Lentiviral Particle

Symbol: CPNE1

Synonyms: COPN1; CPN1

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

**ACCN:** NM\_152925 **ORF Size:** 1611 bp

**ORF Nucleotide** 

- -

Sequence:
OTI Disclaimer:

The ORF insert of this clone is exactly the same as(RC212892).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 152925.2</u>

RefSeq Size: 2022 bp
RefSeq ORF: 1614 bp

Locus ID: 8904

UniProt ID: Q99829

Cytogenetics: 20q11.22

Domains: C2, VWA

**MW:** 59.1 kDa







## **Gene Summary:**

Calcium-dependent membrane-binding proteins may regulate molecular events at the interface of the cell membrane and cytoplasm. This gene encodes a calcium-dependent protein that also contains two N-terminal type II C2 domains and an integrin A domain-like sequence in the C-terminus. However, the encoded protein does not contain a predicted signal sequence or transmembrane domains. This protein has a broad tissue distribution and it may function in membrane trafficking. This gene and the gene for RNA binding motif protein 12 overlap at map location 20q11.21. Alternate splicing results in multiple transcript variants encoding different proteins. [provided by RefSeq, Aug 2008]