

Product datasheet for RC212893L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Liprin alpha 1 (PPFIA1) (NM 177423) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Liprin alpha 1 (PPFIA1) (NM_177423) Human Tagged ORF Clone Lentiviral Particle

Symbol: Liprin alpha 1

Synonyms: LIP.1; LIP1; LIPRIN

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM 177423

ORF Size: 3555 bp

ORF Nucleotide

The ODI

Sequence:
OTI Disclaimer:

The ORF insert of this clone is exactly the same as(RC212893).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 177423.1

 RefSeq Size:
 3945 bp

 RefSeq ORF:
 3558 bp

 Locus ID:
 8500

 UniProt ID:
 Q13136

Cytogenetics: 11q13.3

Protein Families: Druggable Genome, Phosphatase

MW: 133.8 kDa





Gene Summary:

The protein encoded by this gene is a member of the LAR protein-tyrosine phosphatase-interacting protein (liprin) family. Liprins interact with members of LAR family of transmembrane protein tyrosine phosphatases, which are known to be important for axon guidance and mammary gland development. This protein binds to the intracellular membrane-distal phosphatase domain of tyrosine phosphatase LAR, and appears to localize LAR to cell focal adhesions. This interaction may regulate the disassembly of focal adhesion and thus help orchestrate cell-matrix interactions. Alternatively spliced transcript variants encoding distinct isoforms have been described. [provided by RefSeq, Jul 2008]