

## Product datasheet for RC212950L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## CD89 (FCAR) (NM\_133277) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

Product Name: CD89 (FCAR) (NM\_133277) Human Tagged ORF Clone Lentiviral Particle

Symbol: FCAR

Synonyms: CD89; CTB-61M7.2; FcalphaRI

**Mammalian Cell** 

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM 133277

ORF Size: 534 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC212950).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 133277.1

 RefSeq Size:
 1344 bp

 RefSeq ORF:
 537 bp

 Locus ID:
 2204

 UniProt ID:
 P24071

 Cytogenetics:
 19q13.42

**Protein Families:** Transmembrane

**MW:** 19.4 kDa







## **Gene Summary:**

This gene is a member of the immunoglobulin gene superfamily and encodes a receptor for the Fc region of IgA. The receptor is a transmembrane glycoprotein present on the surface of myeloid lineage cells such as neutrophils, monocytes, macrophages, and eosinophils, where it mediates immunologic responses to pathogens. It interacts with IgA-opsonized targets and triggers several immunologic defense processes, including phagocytosis, antibody-dependent cell-mediated cytotoxicity, and stimulation of the release of inflammatory mediators. Multiple alternatively spliced transcript variants encoding different isoforms have been described for this gene. [provided by RefSeq, Jul 2008]