

Product datasheet for RC212960L2V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Beta Arrestin 2 (ARRB2) (NM 199004) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Beta Arrestin 2 (ARRB2) (NM_199004) Human Tagged ORF Clone Lentiviral Particle

Symbol: Beta Arrestin 2

Synonyms: ARB2; ARR2; BARR2

Mammalian Cell

Selection:

None

Vector: pLenti-C-mGFP (PS100071)

Tag: mGFP

ACCN: NM_199004 **ORF Size:** 1182 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC212960).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 199004.1

 RefSeq Size:
 1891 bp

 RefSeq ORF:
 1185 bp

 Locus ID:
 409

 UniProt ID:
 P32121

Cytogenetics: 17p13.2

Protein Families: Druggable Genome

Protein Pathways: Chemokine signaling pathway, Endocytosis, MAPK signaling pathway, Olfactory transduction





Beta Arrestin 2 (ARRB2) (NM_199004) Human Tagged ORF Clone Lentiviral Particle – RC212960L2V

MW: 44.2 kDa

Gene Summary:

Members of arrestin/beta-arrestin protein family are thought to participate in agonist-mediated desensitization of G-protein-coupled receptors and cause specific dampening of cellular responses to stimuli such as hormones, neurotransmitters, or sensory signals. Arrestin beta 2, like arrestin beta 1, was shown to inhibit beta-adrenergic receptor function in vitro. It is expressed at high levels in the central nervous system and may play a role in the regulation of synaptic receptors. Besides the brain, a cDNA for arrestin beta 2 was isolated from thyroid gland, and thus it may also be involved in hormone-specific desensitization of TSH receptors. Multiple alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Mar 2012]