

Product datasheet for RC212993L2V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

FOLR2 (NM_000803) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: FOLR2 (NM_000803) Human Tagged ORF Clone Lentiviral Particle

Symbol: FOLR2

Synonyms: BETA-HFR; FBP; FBP/PL-1; FOLR1; FR-BETA; FR-P3; FRbeta

Mammalian Cell

Selection:

None

Vector: pLenti-C-mGFP (PS100071)

Tag: mGFP

ACCN: NM_000803

ORF Size: 765 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC212993).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 000803.2

 RefSeq Size:
 1119 bp

 RefSeq ORF:
 768 bp

 Locus ID:
 2350

 UniProt ID:
 P14207

 Cytogenetics:
 11q13.4

 Domains:
 Folate_rec

Protein Families: Druggable Genome, Secreted Protein





ORIGENE

MW: 29.1 kDa

Gene Summary:

The protein encoded by this gene is a member of the folate receptor (FOLR) family, and these genes exist in a cluster on chromosome 11. Members of this gene family have a high affinity for folic acid and for several reduced folic acid derivatives, and they mediate delivery of 5-methyltetrahydrofolate to the interior of cells. This protein has a 68% and 79% sequence homology with the FOLR1 and FOLR3 proteins, respectively. Although this protein was originally thought to be specific to placenta, it can also exist in other tissues, and it may play a role in the transport of methotrexate in synovial macrophages in rheumatoid arthritis patients. Multiple transcript variants that encode the same protein have been found for this gene. [provided by RefSeq, Jul 2008]