

## Product datasheet for RC213037L4V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Macrophage inflammatory protein 5 (CCL15) (NM\_032965) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** Macrophage inflammatory protein 5 (CCL15) (NM\_032965) Human Tagged ORF Clone

Lentiviral Particle

Symbol: Macrophage inflammatory protein 5

Synonyms: HCC-2; HMRP-2B; LKN-1; LKN1; MIP-1 delta; MIP-1D; MIP-5; MRP-2B; NCC-3; NCC3; SCYA15;

SCYL3; SY15

**Mammalian Cell** 

Selection:

Puromycin

**Vector:** pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

**ACCN:** NM\_032965

ORF Size: 339 bp

**ORF Nucleotide** 

OTI Disclaimer:

Sequence:

The ORF insert of this clone is exactly the same as(RC213037).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeq:** <u>NM 032965.2</u>

 RefSeq Size:
 955 bp

 RefSeq ORF:
 342 bp

 Locus ID:
 6359

 UniProt ID:
 016663

Cytogenetics: 17q12





## Macrophage inflammatory protein 5 (CCL15) (NM\_032965) Human Tagged ORF Clone Lentiviral Particle - RC213037L4V

Domains: IL8

**Protein Families:** Druggable Genome, Secreted Protein, Transmembrane

**Protein Pathways:** Chemokine signaling pathway, Cytokine-cytokine receptor interaction

MW: 10.1 kDa

**Gene Summary:** This gene is located in a cluster of similar genes in the same region of chromosome 17. These

genes encode CC cytokines, which are secreted proteins characterized by two adjacent cysteines. The product of this gene is chemotactic for T cells and monocytes, and acts through C-C chemokine receptor type 1 (CCR1). The proprotein is further processed into numerous smaller functional peptides. Naturally-occurring readthrough transcripts occur from this gene into the downstream gene, CCL14 (chemokine (C-C motif) ligand 14). [provided

by RefSeq, Jan 2013]