


# Product datasheet for RC213056L4

## Factor XI (F11) (NM\_000128) Human Tagged Lenti ORF Clone

#### **Product data:**

| Product Type:                | Expression Plasmids  |
|------------------------------|--|
| Product Name:                | Factor XI (F11) (NM_000128) Human Tagged Lenti ORF Clone                               |
| Tag:                         | mGFP   |
| Symbol:                      | Factor XI  |
| Synonyms:                    | FXI; PTA   |
| Mammalian Cell<br>Selection: | Puromycin  |
| Vector:                      | pLenti-C-mGFP-P2A-Puro (PS100093)  |
| E. coli Selection:           | Chloramphenicol (34 ug/mL)   |
| ORF Nucleotide<br>Sequence:  | The ORF insert of this clone is exactly the same as(RC213056).                         |
| <b>Restriction Sites:</b>    | Sgfl-Mlul  |
| Cloning Scheme:              |  |
|                              | Cloning sites used for ORF Shuttling:  |
|                              | Sgf I         ORF         Mlu I          GCG ATC GCC         ATG// NNN         ACG CGT |



ACCN: ORF Size: NM\_000128 1875 bp

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


| <b>GRIGENE</b> Factor  | XI (F11) (NM_000128) Human Tagged Lenti ORF Clone – RC213056L4  |
|------------------------|---|
| OTI Disclaimer:        | The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>   |
| OTI Annotation:        | This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.  |
| Components:            | The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).  |
| Reconstitution Method: | <ol> <li>Centrifuge at 5,000xg for 5min.</li> <li>Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li> <li>Close the tube and incubate for 10 minutes at room temperature.</li> <li>Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li> <li>Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li> </ol>  |
| RefSeq:                | <u>NM 000128.2</u>  |
| RefSeq Size:           | 2217 bp   |
| RefSeq ORF:            | 1878 bp   |
| Locus ID:              | 2160  |
| UniProt ID:            | <u>P03951</u>   |
| Cytogenetics:          | 4q35.2  |
| Domains:               | APPLE, Tryp_SPc, PAN  |
| Protein Families:      | Druggable Genome, Protease, Secreted Protein  |
| Protein Pathways:      | Complement and coagulation cascades   |
| MW:                    | 69.9 kDa  |
| Gene Summary:          | This gene encodes coagulation factor XI of the blood coagulation cascade. This protein is present in plasma as a zymogen, which is a unique plasma coagulation enzyme because it exists as a homodimer consisting of two identical polypeptide chains linked by disulfide bonds. During activation of the plasma factor XI, an internal peptide bond is cleaved by factor XIIa (or XII) in each of the two chains, resulting in activated factor XIa, a serine protease composed of two heavy and two light chains held together by disulfide bonds. This activated plasma factor XI triggers the middle phase of the intrisic pathway of blood coagulation by activating factor IX. Defects in this factor lead to Rosenthal syndrome, a blood coagulation |

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


abnormality. [provided by RefSeq, Jul 2008]



### **Product images:**



Circular map for RC213056L4



Double digestion of RC213056L4 using Sgfl and Mlul

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US