

Product datasheet for RC213065L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Dopamine D2 Receptor (DRD2) (NM_016574) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Dopamine D2 Receptor (DRD2) (NM_016574) Human Tagged ORF Clone Lentiviral Particle

Symbol: Dopamine D2 Receptor

Synonyms: D2DR; D2R

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_016574 **ORF Size:** 1242 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC213065).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 016574.2

 RefSeq Size:
 2556 bp

 RefSeq ORF:
 1245 bp

 Locus ID:
 1813

 UniProt ID:
 P14416

 Cytogenetics:
 11q23.2

Protein Families: Druggable Genome, GPCR, Transmembrane

Protein Pathways: Gap junction, Neuroactive ligand-receptor interaction





Dopamine D2 Receptor (DRD2) (NM_016574) Human Tagged ORF Clone Lentiviral Particle -RC213065L4V

MW: 47.2 kDa

Gene Summary:

This gene encodes the D2 subtype of the dopamine receptor. This G-protein coupled receptor inhibits adenylyl cyclase activity. A missense mutation in this gene causes myoclonus dystonia; other mutations have been associated with schizophrenia. Alternative splicing of this gene results in two transcript variants encoding different isoforms. A third variant has been described, but it has not been determined whether this form is normal or due to aberrant splicing. [provided by RefSeq, Jul 2008]