

Product datasheet for RC213128

MAX (NM_002382) Human Tagged ORF Clone

Product data:

Product Type: Expression Plasmids
 Product Name: MAX (NM_002382) Human Tagged ORF Clone
 Tag: Myc-DDK
 Symbol: MAX
 Synonyms: bHLHd4
 Mammalian Cell Selection: Neomycin
 Vector: pCMV6-Entry (PS100001)
 E. coli Selection: Kanamycin (25 ug/mL)
 ORF Nucleotide Sequence: >RC213128 representing NM_002382
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGATCGCC**

ATGAGCGATAACGATGACATCGAGGTGGAGAGCGACGAAGAGCAACCGAGTTTCAATCTGCGGCTGACA
 AACGGGCTCATCATAATGCACTGGAACGAAAACGTAGGGACCACATCAAAGACAGCTTTCACAGTTTGGC
 GGACTCAGTCCCATCACTCCAAGGAGAGAAGGCATCCCGGGCCAAATCCTAGACAAAGCCACAGAATAT
 ATCCAGTATATGCGAAGGAAAAACACACACACCAGCAAGATATTGACGACCTCAAGCGGCAGAATGCTC
 TTCTGGAGCAGCAAGTCCGTGCACTGGAGAAGGCGAGGTCAAGTGCCCACTGCAGACCAACTACCCCTC
 CTAGACAACAGCCTCTACACCAAGCCAAAGGGCAGCACCATCTCTGCCTTCGATGGGGGCTCGGACTCC
 AGCTCGGAGTCTGAGCCTGAAGAGCCCCAAAGCAGGAAGAAGCTCCGGATGGAGGCCAGC

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >RC213128 representing NM_002382
 Red=Cloning site Green=Tags(s)

MSDNDIEVESDEEQPRFQSAADKRAHHNALERKRRDHKDSFHSLRDSVPSLQGEKASRAQILDKATEY
 IQYMRKNHHTHQDIDDLKRNALLEQQVRALEKARSSAQLQTNYPSSDNLTYNAKGSTISAFDGGSDS
 SSESEPEEPQSRKCLRMEAS

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites: Sgfl-MluI



[View online >](#)

Cloning Scheme:

ACCN:

NM_002382

ORF Size:

480 bp

OTI Disclaimer:

Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation:

This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components:

The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).


Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_002382.5](#)
RefSeq Size: 2068 bp
RefSeq ORF: 483 bp
Locus ID: 4149
UniProt ID: [P61244](#)
Cytogenetics: 14q23.3
Domains: HLH
Protein Families: Druggable Genome, Transcription Factors
Protein Pathways: MAPK signaling pathway, Pathways in cancer, Small cell lung cancer
MW: 18.1 kDa

Gene Summary: The protein encoded by this gene is a member of the basic helix-loop-helix leucine zipper (bHLHZ) family of transcription factors. It is able to form homodimers and heterodimers with other family members, which include Mad, Mxi1 and Myc. Myc is an oncoprotein implicated in cell proliferation, differentiation and apoptosis. The homodimers and heterodimers compete for a common DNA target site (the E box) and rearrangement among these dimer forms provides a complex system of transcriptional regulation. Mutations of this gene have been reported to be associated with hereditary pheochromocytoma. A pseudogene of this gene is located on the long arm of chromosome 7. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Aug 2012]

Product images:



Circular map for RC213128