

# Product datasheet for RC213165L2V

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## CD8A (NM\_171827) Human Tagged ORF Clone Lentiviral Particle

#### **Product data:**

**Product Type:** Lentiviral Particles

**Product Name:** CD8A (NM\_171827) Human Tagged ORF Clone Lentiviral Particle

Symbol: CD8A

Synonyms: CD8; Leu2; p32

Mammalian Cell

Selection:

None

**Vector:** pLenti-C-mGFP (PS100071)

Tag: mGFP

**ACCN:** NM\_171827

ORF Size: 594 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC213165).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 171827.1</u>

RefSeq Size: 2150 bp
RefSeq ORF: 597 bp
Locus ID: 925

 UniProt ID:
 P01732

 Cytogenetics:
 2p11.2

Protein Families: Adult stem cells, Druggable Genome, ES Cell Differentiation/IPS, Secreted Protein,

Transmembrane





### CD8A (NM\_171827) Human Tagged ORF Clone Lentiviral Particle - RC213165L2V

Protein Pathways: Antigen processing and presentation, Cell adhesion molecules (CAMs), Hematopoietic cell

lineage, Primary immunodeficiency, T cell receptor signaling pathway

**MW:** 19.4 kDa

**Gene Summary:** The CD8 antigen is a cell surface glycoprotein found on most cytotoxic T lymphocytes that

mediates efficient cell-cell interactions within the immune system. The CD8 antigen acts as a coreceptor with the T-cell receptor on the T lymphocyte to recognize antigens displayed by an antigen presenting cell in the context of class I MHC molecules. The coreceptor functions as either a homodimer composed of two alpha chains or as a heterodimer composed of one alpha and one beta chain. Both alpha and beta chains share significant homology to immunoglobulin variable light chains. This gene encodes the CD8 alpha chain. Multiple transcript variants encoding different isoforms have been found for this gene. The major protein isoforms of this gene differ by the presence or absence of a transmembrane domain and thus differ in being a membrane-anchored or secreted protein. [provided by RefSeq, May

2020]