

Product datasheet for RC213184L2V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

HMGCR (NM_000859) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: HMGCR (NM_000859) Human Tagged ORF Clone Lentiviral Particle

Symbol: HMGCR Synonyms: LDLCQ3

Mammalian Cell None

Selection:

Vector:

pLenti-C-mGFP (PS100071)

Tag: mGFP

ACCN: NM_000859 **ORF Size:** 2664 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC213184).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 000859.1

 RefSeq Size:
 4471 bp

 RefSeq ORF:
 2667 bp

 Locus ID:
 3156

 UniProt ID:
 P04035

 Cytogenetics:
 5q13.3

Domains: HMG-CoA red

Protein Families: Druggable Genome, Transmembrane





HMGCR (NM_000859) Human Tagged ORF Clone Lentiviral Particle - RC213184L2V

Protein Pathways: Metabolic pathways, Terpenoid backbone biosynthesis

MW: 97.3 kDa

Gene Summary: HMG-CoA reductase is the rate-limiting enzyme for cholesterol synthesis and is regulated via

a negative feedback mechanism mediated by sterols and non-sterol metabolites derived from mevalonate, the product of the reaction catalyzed by reductase. Normally in mammalian cells this enzyme is suppressed by cholesterol derived from the internalization and degradation of low density lipoprotein (LDL) via the LDL receptor. Competitive inhibitors of the reductase induce the expression of LDL receptors in the liver, which in turn increases the catabolism of plasma LDL and lowers the plasma concentration of cholesterol, an important determinant of atherosclerosis. Alternatively spliced transcript variants encoding different isoforms have

been found for this gene. [provided by RefSeq, Aug 2008]