

OriGene Technologies, Inc.


9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for RC213206L2

Eph receptor A5 (EPHA5) (NM_004439) Human Tagged Lenti ORF Clone

Product data:

| Product Type: | Expression Plasmids |
|------------------------------|---|
| Product Name: | Eph receptor A5 (EPHA5) (NM_004439) Human Tagged Lenti ORF Clone |
| Tag: | mGFP |
| Symbol: | Eph receptor A5 |
| Synonyms: | CEK7; EHK-1; EHK1; EK7; HEK7; TYRO4 |
| Mammalian Cell Selection: | None |
| Vector: | pLenti-C-mGFP (PS100071) |
| E. coli Selection: | Chloramphenicol (34 ug/mL) |
| ORF Nucleotide Sequence: | The ORF insert of this clone is exactly the same as(RC213206). |
| Restriction Sites: | Sgfl-Mlul |
| Cloning Scheme: | |
| | Cloning sites used for ORF Shuttling: Sgf 1 ORF Mlu 1 GCG ATC GCC ATG// NNN ACG CGT |




* The last codon before the Stop codon of the ORF.



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Plasmid Map:



| ACCN: | NM_004439 |
|-----------------|--|
| ORF Size: | 3111 bp |
| OTI Disclaimer: | Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at <u>custsupport@origene.com</u> or by calling 301.340.3188 option 3 for pricing and delivery. |
| | The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u> |
| OTI Annotation: | This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene. |
| Components: | The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water). |

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

PORIGENE Eph receptor A5 (EPHA5) (NM_004439) Human Tagged Lenti ORF Clone – RC213206L2

| Reconstitution Method: | Centrifuge at 5,000xg for 5min. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. Close the tube and incubate for 10 minutes at room temperature. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C. |
|------------------------|---|
| RefSeq: | <u>NM 004439.4, NP 004430.3</u> |
| RefSeq Size: | 3349 bp |
| RefSeq ORF: | 3114 bp |
| Locus ID: | 2044 |
| UniProt ID: | <u>P54756</u> |
| Cytogenetics: | 4q13.1-q13.2 |
| Domains: | pkinase, EPH_lbd, TyrKc, SAM, S_TKc, FN3 |
| Protein Families: | Druggable Genome, Protein Kinase, Transmembrane |
| Protein Pathways: | Axon guidance |
| MW: | 114.78 kDa |
| Gene Summary: | This gene belongs to the ephrin receptor subfamily of the protein-tyrosine kinase family. EPH and EPH-related receptors have been implicated in mediating developmental events, particularly in the nervous system. Receptors in the EPH subfamily typically have a single kinase domain and an extracellular region containing a Cys-rich domain and 2 fibronectin type III repeats. The ephrin receptors are divided into 2 groups based on the similarity of their extracellular domain sequences and their affinities for binding ephrin-A and ephrin-B ligands. Alternatively spliced transcript variants encoding different isoforms have been described. |

[provided by RefSeq, Aug 2013]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US