

Product datasheet for **RC213207L4V**

Ikaros (IKZF1) (NM_006060) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Ikaros (IKZF1) (NM_006060) Human Tagged ORF Clone Lentiviral Particle
Symbol:	Ikaros
Synonyms:	CVID13; Hs.54452; IK1; IKAROS; LyF-1; LYF1; PPP1R92; PRO0758; ZNFN1A1
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_006060
ORF Size:	1557 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC213207).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	NM_006060.2
RefSeq Size:	3962 bp
RefSeq ORF:	1560 bp
Locus ID:	10320
UniProt ID:	Q13422
Cytogenetics:	7p12.2
Domains:	zf-C2H2
Protein Families:	Druggable Genome, Transcription Factors




[View online »](#)

MW: 57.3 kDa

Gene Summary: This gene encodes a transcription factor that belongs to the family of zinc-finger DNA-binding proteins associated with chromatin remodeling. The expression of this protein is restricted to the fetal and adult hemo-lymphopoietic system, and it functions as a regulator of lymphocyte differentiation. Several alternatively spliced transcript variants encoding different isoforms have been described for this gene. Most isoforms share a common C-terminal domain, which contains two zinc finger motifs that are required for hetero- or homo-dimerization, and for interactions with other proteins. The isoforms, however, differ in the number of N-terminal zinc finger motifs that bind DNA and in nuclear localization signal presence, resulting in members with and without DNA-binding properties. Only a few isoforms contain the requisite three or more N-terminal zinc motifs that confer high affinity binding to a specific core DNA sequence element in the promoters of target genes. The non-DNA-binding isoforms are largely found in the cytoplasm, and are thought to function as dominant-negative factors. Overexpression of some dominant-negative isoforms have been associated with B-cell malignancies, such as acute lymphoblastic leukemia (ALL). [provided by RefSeq, May 2014]

Product images:



[RC213207L4] was used to prepare Lentiviral particles using [TR30037] packaging kit. HEK293T cells were transduced with RC213207L4V particle to overexpress human IKZF1-mGFP fusion protein.