

## Product datasheet for **RC213342**

### **THRAP3 (NM\_005119) Human Tagged ORF Clone**

#### **Product data:**

Product Type:	Expression Plasmids
Product Name:	THRAP3 (NM_005119) Human Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	THRAP3
Synonyms:	BCLAF2; TRAP150
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)

**ORF Nucleotide Sequence:**

>RC213342 ORF sequence  
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
 GCC**CGCATCGCC**

ATGTCAAAAACAACAATCCAAGTCTGGATCTCGCTCTTCTCGCTCAAGATCTGCATCAAGATCTCGTT  
 CTGTTTCAATTTTCGAAGTCTCGGTCCCGAAGCCGATCTCTCTCGTTCAAGGAAGCGCAGGCTGAGTTC  
 TAGGTCTCGTTCCAGATCATATTCTCCAGCTCATAACAGAGAAAAGAAACCCCAAGAGTATATCAGAAT  
 CGGGATTTCCGAGGTCAACAACAGAGGCTATAGAAGGCCCTATTATTTCCGTGGGCGTAACAGAGGCTTTT  
 ATCCATGGGGCAATATAACCGAGGAGGCTATGGAACTACCGCTCAAATGGCAGAATTACCGCAAGC  
 ATACAGTCTCGTCGAGGCCGTTCAAGATCCCGGTCCCAAGAGAAGGTCCCCTTACCAAGGTCCAGG  
 AGCCATTTAGAACTCTGATAAGTCGTCTTCTGACCGGTCAAGGCGCTCCTCATCTCCGTTCTTCTCT  
 CCAACCATAGCCGAGTTGAATCTTCTAAGCGCAAGTCTGCAAAGGAGAAAAAGTCTCTTCTAAGGATAG  
 CCGGCCATCTCAGGCTGCCGGGATAACCAGGGAGATGAGCCAAGGAGCAGACATTTCTCGGAGGCACC  
 TCTCAAGATACAAAAGCATCTGAGAGCTCGAAGCCATGGCCAGATGCCACCTACGGCACTGGTTCTGCAT  
 CACGGGCTCAGCAGTTTCTGAGCTGAGTCTCGGGAGCGAAGCCAGCTCTCAAAAGCCCCCTCCAGTC  
 TGTGGTGGTGAAGCGCGGTACCCCGTCTAGCCCGTGCCAAAACCTAGTCTCCACTTTCCAGCACA  
 TCCCAGATGGGCTCAACTCTGCCGAGTGGTGCCGGGATCAGTCTGGGACACCAAGGTGAGTTCGACC  
 ATGGTTCTGGGTCCCTGAGTCCATCCAAAAGAGCCCTGTGGGTAAAGTCCACCATCCACTGGCTCCAC  
 ATATGGCTCATCTCAGAAGGAGGAGTGTCTGTTTCCAGGAGGAGCAGCCTATACAAAGAGGTATCTAGAA  
 GAGCAGAAGACAGAGAATGGAAAAGATAAGGAACAGAAAACAACAATACCGATAAAGAAAAAATAAAG  
 AGAAAGGAGCTTCTCTGACACAGGCTTGGGTGATGGAAAAATGAAATCTGATTCTTTTGTCTCCAAAAC  
 TGATTCTGAGAAGCCTTTTCGGGGCAGTCAAGTCTCCAAAAGGTATAAGCTCCGAGATGACTTTGAGAAG  
 AAGATGGCTGACTTCCACAAGGAGGAGATGGATGATCAAGATAAGGACAAAGCTAAGGGAAGAAAGGAAT  
 CTGAGTTTGTGATGAACCCAAATTTATGTCTAAAGTCAAGGTGCAAACAAAACAGGAGGAGGAGAA  
 GTCAGGCAAAATGGGAGGGCTGGTATATGCACCTCCAGGGAAGGAAAAGCAGAGAAAAACAGAGGAGCTG  
 GAGGAGGAGTCTTTCCAGAGAGATCCAAAAGGAAGTCCGGGCAAGAGAAGCGAAGGTGGGCACAGGG  
 GCTTTGTGCTGAGAAGAATTTCCGAGTACTGCTTATAAAGCAGTCCAGGAGAAAAGCTCATCACCTCC  
 CCCAAGAAAGACCTCTGAGAGCCGAGACAAGTGGGAGCGAAAGGAGATTTCCACAGGAAAGTCTTCC  
 TTTTCCATTACTCGAGAGGCACAGTCAATGTCCGGATGGACTCTTTTGTGAGGACCTCGCACGCCCA  
 GTGGCTTATTGGCTCAGGAACGCAAGCTTTGCCGAGATCTAGTCCATAGCAACAAAAGGAACAGGAGTT  
 TCGTTCCATTTTCCAGCACATACATCAGCTCAGTCTCAGCGTAGCCCTCAGAAGTGTTTGCCAACAT  
 ATAGTGACCATTGTTACCATGTTAAAGAGCATCACTTTGGGTCTCAGGAATGACATTACATGAACGCT  
 TTAATAATACCTAAAGAGAGGAAGTGAAGCAGGAGGCAGCCAAAACAAGAAAAGCCAGAGATACACAG  
 GAGAATAGACATTTCCCCAGTACATTCAGAAAACATGGTTTGGCTCATGATGAAATGAAAAGTCCCCGG  
 GAACCTGGCTACAAGGCTGAGGGAAAATACAAAGATGATCCTGTTGATCTCCGCTTGATATTGAACGTC  
 GTAATAAACATAAGGAGAGAGATCTTAAACGAGGTAATCGAGAGAATCAGTGGATCCCGAGACTCCAG  
 TCACTCAAGGGAAGGTGAGTCAAAAAACAGAGAAAACCTATAAAGGATCAAAGAAAACAGAAAGCAT  
 CGGAGAGCAAGAGACAGGTCCAGATCCTCCTCTTCCCTCCAGTCACTCACTCTACAAAGCAGAAG  
 AGTACACTGAAGAGACAGAGGAAAGAGAGGAGAGCACCAGGGCTTTGACAAATCAAGACTGGGGACCAA  
 AGACTTTGTGGTCCAAGTGAAAGAGGAGGTGGCAGAGCTCGAGGAACCTTTTCAAGTTTCCAGCCAGAGGA  
 AGAGGCTGGGGCAGAGGCAACTACTCTGGGAACAATAACAACAACAGCAACAACGATTTTCAAAAAAGAA  
 ACCGGGAAGAGGAGTGGGACCCAGAGTACACCCAAAAGCAAGAAGTATTACTTGCATGATGACCGTGA  
 AGGCGAAGGCAGTGACAAGTGGGTGAGCCGGGGCCGGGGCCGAGGAGCCTTCTCGGGGTGGGGCCGG  
 TTCATGTTCCGGAATCAAGTACCAGCCCAAGTGGGCCATGACAAGTTCAGTGGGAGGAAGGGGAGA  
 TTGAAGACGACGAGAGTGGGACAGAGAACCAGAAAGAGAAGGACAATATACAGCCCAACACCGAG

**ACGCGT**ACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT  
 ACAAGGATGACGACGATAAGGTTTAA

**Protein Sequence:** >RC213342 protein sequence  
Red=Cloning site Green=Tags(s)

```
MSKTNKSKSGSRSSRSASRSRSLRSLRKRRLSSRSRSYSPAHRNRNHPRVYQN
RDFRGHNRYRRPYFRGRNRFYPWGQYNRGGYGNYSNWQNYRQAYSRRGRSRSPKRRSPRSR
SHSRNSDKSSSDRSRRSSSSRNHSRVESKRKSAKEKSSSKDSRPSQAAGDNQGDQTFSGGT
SQDTKASESSKPWPDATYGTGSASRASAVSELSPRERSPALQSPLQSVVRRRSPRSPVPKPSPLSST
SQMGSTLPSGAGYQSGTHQGFQDHGSGSLSPSKSPVKGSPSTGSTYGSQKEESAASGGAAYTKRYLE
EQKTENGKDKQQTNTDKEKIKEKGSFSDTGLGDGKMKSDSFAPKTDSEKPFRRGSPKRYKLDDFEK
KMADFHKEEMDDQDKAKGRKESEFDDEPKFMSKIVIGANKNQEESKSGWEGLVYAPPGKEKQRKTEEL
EEESFPERSKEDRGKRSEGGHRGFVPEKNFRVTAYKAVQEKSSSPPRKTSESRDKLGAKGDFPTGKSS
FSITREAQVNRMDSFDEDLARPSGLLAQERKLCRDLVHSNKKEQEFRSIFQHIQSAQSQRSPSELFAQH
IVTIVHHVKEHHFGSSGMLHERFTKYLKRGTEQEAANKKSPEIHRRIDISPSTFRKHGLAHDEMKSPPR
EPGYKAEGKYKDDPVDLRDIERRKKHKERDLKRGKSRESVDSRDSHRSRERSAEKTEKTHKSKKQKKH
RRARDRSSSSSSQSSHSYKAEYETEEREESTTGFDKSRGKDFVGPSEGGGRARGTFFQFRARG
RGWGRGNYSNNNNNSNDFQKRNREEEDPEYTPKSKYYLHDDREGESDKWVSRGRGRGAFPRGRGR
FMFRKSSTSPKWAHDKFSGEEGIEDDESGTENREEKDNIQPTTE
```


TRTRPLEQKLISEEDLAANDILDYKDDDDKV

**Chromatograms:** [https://cdn.origene.com/chromatograms/mk6602\\_f07.zip](https://cdn.origene.com/chromatograms/mk6602_f07.zip)

**Restriction Sites:** SgfI-MluI

**Cloning Scheme:**

Cloning sites used for ORF Shuttling:



\* The last codon before the Stop codon of the ORF


**ACCN:** NM\_005119

**ORF Size:** 2865 bp


<b>OTI Disclaimer:</b>	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
<b>OTI Annotation:</b>	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>Note:</b>	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
<b>RefSeq:</b>	<a href="#">NM_005119.4</a>
<b>RefSeq Size:</b>	4447 bp
<b>RefSeq ORF:</b>	2868 bp
<b>Locus ID:</b>	9967
<b>UniProt ID:</b>	<a href="#">Q9Y2W1</a>
<b>Cytogenetics:</b>	1p34.3
<b>Protein Families:</b>	Druggable Genome, Transcription Factors
<b>MW:</b>	108.7 kDa

**Gene Summary:**

Involved in pre-mRNA splicing. Remains associated with spliced mRNA after splicing which probably involves interactions with the exon junction complex (EJC). Can trigger mRNA decay which seems to be independent of nonsense-mediated decay involving premature stop codons (PTC) recognition. May be involved in nuclear mRNA decay. Involved in regulation of signal-induced alternative splicing. During splicing of PTPRC/CD45 is proposed to sequester phosphorylated SFQ from PTPRC/CD45 pre-mRNA in resting T-cells. Involved in cyclin-D1/CCND1 mRNA stability probably by acting as component of the SNARP complex which associates with both the 3'end of the CCND1 gene and its mRNA. Involved in response to DNA damage. Is excluded from DNA damage sites in a manner that parallels transcription inhibition; the function may involve the SNARP complex. Initially thought to play a role in transcriptional coactivation through its association with the TRAP complex; however, it is not regarded as a stable Mediator complex subunit. Cooperatively with HELZ2, enhances the transcriptional activation mediated by PPARG, maybe through the stabilization of the PPARG binding to DNA in presence of ligand. May play a role in the terminal stage of adipocyte differentiation. Plays a role in the positive regulation of the circadian clock. Acts as a coactivator of the CLOCK-ARNTL/BMAL1 heterodimer and promotes its transcriptional activator activity and binding to circadian target genes (PubMed:24043798).[UniProtKB/Swiss-Prot Function]

**Product images:**


Circular map for RC213342



Western blot validation of overexpression lysate (Cat# [LY417509]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC213342 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).