

## Product datasheet for RC213377L4V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## IL17RC (NM\_153460) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** IL17RC (NM\_153460) Human Tagged ORF Clone Lentiviral Particle

Symbol: IL17RC

Synonyms: CANDF9; IL17-RL; IL17RL

**Mammalian Cell** 

Selection:

Puromycin

**Vector:** pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

**ACCN:** NM\_153460 **ORF Size:** 2160 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC213377).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 153460.1, NP 703190.1

 RefSeq Size:
 2478 bp

 RefSeq ORF:
 2163 bp

 Locus ID:
 84818

 UniProt ID:
 Q8NAC3

Cytogenetics: 3p25.3-p24.1

**Protein Families:** Druggable Genome, Transmembrane

**MW:** 76.4 kDa







## **Gene Summary:**

This gene encodes a single-pass type I membrane protein that shares similarity with the interleukin-17 receptor (IL-17RA). Unlike IL-17RA, which is predominantly expressed in hemopoietic cells, and binds with high affinity to only IL-17A, this protein is expressed in nonhemopoietic tissues, and binds both IL-17A and IL-17F with similar affinities. The proinflammatory cytokines, IL-17A and IL-17F, have been implicated in the progression of inflammatory and autoimmune diseases. Multiple alternatively spliced transcript variants encoding different isoforms have been detected for this gene, and it has been proposed that soluble, secreted proteins lacking transmembrane and intracellular domains may function as extracellular antagonists to cytokine signaling. [provided by RefSeq, Feb 2011]