

## Product datasheet for RC213525L1V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## RASSF1 (NM 007182) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type: Lentiviral Particles** 

**Product Name:** RASSF1 (NM\_007182) Human Tagged ORF Clone Lentiviral Particle

Symbol:

123F2; NORE2A; RASSF1A; RDA32; REH3P21 Synonyms:

**Mammalian Cell** 

Selection:

None

Vector: pLenti-C-Myc-DDK (PS100064)

Myc-DDK Tag: NM 007182 ACCN: **ORF Size:** 1020 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC213525).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 007182.4

RefSeq Size: 1968 bp RefSeq ORF: 1023 bp Locus ID: 11186 **UniProt ID:** Q9NS23 Cytogenetics: 3p21.31

**Domains:** RA, DAG\_PE-bind

**Protein Families:** Druggable Genome





## RASSF1 (NM\_007182) Human Tagged ORF Clone Lentiviral Particle - RC213525L1V

**Protein Pathways:** Bladder cancer, Non-small cell lung cancer, Pathways in cancer

MW: 38.6 kDa

**Gene Summary:** This gene encodes a protein similar to the RAS effector proteins. Loss or altered expression

of this gene has been associated with the pathogenesis of a variety of cancers, which

suggests the tumor suppressor function of this gene. The inactivation of this gene was found to be correlated with the hypermethylation of its CpG-island promoter region. The encoded protein was found to interact with DNA repair protein XPA. The protein was also shown to inhibit the accumulation of cyclin D1, and thus induce cell cycle arrest. Several alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported.

[provided by RefSeq, May 2011]