

Product datasheet for RC213525L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

RASSF1 (NM_007182) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: RASSF1 (NM_007182) Human Tagged ORF Clone Lentiviral Particle

Symbol: RASSF1

Synonyms: 123F2; NORE2A; RASSF1A; RDA32; REH3P21

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

 Tag:
 Myc-DDK

 ACCN:
 NM_007182

 ORF Size:
 1020 bp

ORF Nucleotide

OTI Disclaimer:

.

Sequence:

The ORF insert of this clone is exactly the same as(RC213525).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 007182.4

 RefSeq Size:
 1968 bp

 RefSeq ORF:
 1023 bp

 Locus ID:
 11186

 UniProt ID:
 Q9NS23

Cytogenetics: 3p21.31

Domains: RA, DAG_PE-bind
Protein Families: Druggable Genome





RASSF1 (NM_007182) Human Tagged ORF Clone Lentiviral Particle - RC213525L3V

Protein Pathways: Bladder cancer, Non-small cell lung cancer, Pathways in cancer

MW: 38.6 kDa

Gene Summary: This gene encodes a protein similar to the RAS effector proteins. Loss or altered expression

of this gene has been associated with the pathogenesis of a variety of cancers, which suggests the tumor suppressor function of this gene. The inactivation of this gene was found to be correlated with the hypermethylation of its CpG-island promoter region. The encoded

protein was found to interact with DNA repair protein XPA. The protein was also shown to inhibit the accumulation of cyclin D1, and thus induce cell cycle arrest. Several alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported.

[provided by RefSeq, May 2011]