

Product datasheet for RC213540L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

SCF (KITLG) (NM_003994) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: SCF (KITLG) (NM_003994) Human Tagged ORF Clone Lentiviral Particle

Symbol: SCF

Synonyms: DCUA; DFNA69; FPH2; FPHH; Kitl; KL-1; MGF; SCF; SF; SHEP7; SLF

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM 003994

ORF Size: 735 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC213540).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 003994.4</u>

 RefSeq Size:
 5351 bp

 RefSeq ORF:
 738 bp

 Locus ID:
 4254

 UniProt ID:
 P21583

 Cytogenetics:
 12q21.32

Domains: SCF

Protein Families: Druggable Genome, Transmembrane





SCF (KITLG) (NM_003994) Human Tagged ORF Clone Lentiviral Particle - RC213540L3V

Protein Pathways: Cytokine-cytokine receptor interaction, Hematopoietic cell lineage, Melanogenesis, Pathways

in cancer

MW: 27.87 kDa

Gene Summary: This gene encodes the ligand of the tyrosine-kinase receptor encoded by the KIT locus. This

ligand is a pleiotropic factor that acts in utero in germ cell and neural cell development, and

hematopoiesis, all believed to reflect a role in cell migration. In adults, it functions

pleiotropically, while mostly noted for its continued requirement in hematopoiesis. Two transcript variants encoding different isoforms have been found for this gene. [provided by

RefSeq, Jul 2008]