

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for RC213609L4V

ICK (CILK1) (NM_014920) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	ICK (CILK1) (NM_014920) Human Tagged ORF Clone Lentiviral Particle
Symbol:	CILK1
Synonyms:	ECO; EJM10; hICK; ICK; LCK2; MRK
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_014920
ORF Size:	1896 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC213609).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 014920.2</u>
RefSeq Size:	6095 bp
RefSeq ORF:	1899 bp
Locus ID:	22858
UniProt ID:	<u>Q9UPZ9</u>
Cytogenetics:	6p12.1
Protein Families:	Druggable Genome, Protein Kinase
MW:	71.2 kDa



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Gene Summary:Eukaryotic protein kinases are enzymes that belong to a very extensive family of proteins
which share a conserved catalytic core common with both serine/threonine and tyrosine
protein kinases. This gene encodes an intestinal serine/threonine kinase harboring a dual
phosphorylation site found in mitogen-activating protein (MAP) kinases. The protein localizes
to the intestinal crypt region and is thought to be important in intestinal epithelial cell
proliferation and differentiation. Alternative splicing has been observed at this locus and two
variants, encoding the same isoform, have been identified. [provided by RefSeq, Jul 2008]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US