

Product datasheet for RC213689L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Eph receptor A1 (EPHA1) (NM 005232) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Eph receptor A1 (EPHA1) (NM_005232) Human Tagged ORF Clone Lentiviral Particle

Symbol: Eph receptor A1
Synonyms: EPH; EPHT; EPHT1

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

 Tag:
 Myc-DDK

 ACCN:
 NM_005232

 ORF Size:
 2928 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC213689).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 005232.3

 RefSeq Size:
 3367 bp

 RefSeq ORF:
 2931 bp

 Locus ID:
 2041

 UniProt ID:
 P21709

 Cytogenetics:
 7q34-q35

Protein Families: Druggable Genome, Protein Kinase, Transmembrane

Protein Pathways: Axon guidance





Eph receptor A1 (EPHA1) (NM_005232) Human Tagged ORF Clone Lentiviral Particle – RC213689L3V

MW: 107.9 kDa

Gene Summary:

This gene belongs to the ephrin receptor subfamily of the protein-tyrosine kinase family. EPH and EPH-related receptors have been implicated in mediating developmental events, particularly in the nervous system. Receptors in the EPH subfamily typically have a single kinase domain and an extracellular region containing a Cys-rich domain and 2 fibronectin type III repeats. The ephrin receptors are divided into 2 groups based on the similarity of their extracellular domain sequences and their affinities for binding ephrin-A and ephrin-B ligands. This gene is expressed in some human cancer cell lines and has been implicated in carcinogenesis. [provided by RefSeq, Jul 2008]