

Product datasheet for RC213905

Constitutive androstane receptor (NR1I3) (NM_001077482) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Constitutive androstane receptor (NR1I3) (NM_001077482) Human Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	Constitutive androstane receptor
Synonyms:	CAR; CAR1; MB67
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
ORF Nucleotide Sequence:	>RC213905 representing NM_001077482 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGGCCAGTAGGGAAGATGAGCTGAGGAAGTGTGTGGTATGTGGGGACCAAGCCACAGGCTACCACTTTA
ATGCGCTGACTTGTGAGGGCTGCAAGGGTTTCTCAGGAGAACAGTCAGCAAAGCATTGGTCCCACCTG
CCCCTTTGCTGGAAGCTGTGAAGTCAGCAAGACTCAGAGGCGCCACTGCCAGCCTGCAGGTTGCAGAAG
TGCTTAGATGCTGGCATGAGGAAAGACATGATACTGTCCGAGCAAGCCCTGGCATTGCGGCGAGCAAAGC
AGGCCAGCGGGCCACAGCAAACACCTGTGCAACTGAGTAAGGAGCAAGAAGAGCTGATCCGGACT
CCTGGGGCCACACCCGCCACATGGGCACCATGTTTGAACAGTTTGTGCAGTTTAGGCCTCCAGCTCAT
CTGTTTCATCCATCACCAGCCCTTGCCACCCCTGGCCCTGTGCTGCCTCTGGTCACACACTTCGAGACA
TCAACTTTTCATGGTACTGCAAGTCACTCAAGTTTACTAAGGACCTGCCGTCTCCGTTCCCTGCCAT
TGAAGACCAGATCTCCCTTCTCAAGGGAGCAGCTGTGAAATCTGTCACATCGTACTCAATACCACTTTC
TGTCTCCAACACAAAATTCTCTGCGGGCTCTTCGCTACACAATTGAAGATGGAGCCCGTGTATCTC
CCACAGTGGGGTCCAGGTAGAGTTTTTGGAGTTGCTCTTTCCTTCCATGGAACACTACGAAAAGTCA
GCTCCAAGAGCCTGAGTATGTGCTCTTGGCTGCCATGGCCCTCTCTCTCTGCTCCCTATCTTACAGAC
CGACCTGGAGTTACCCAGAGAGATGAGATTGATCAGCTGCAAGAGGAGATGGCACTGACTGCAAAAGCT
ACATCAAGGGCCAGCAGCGAAGGCCCGGGATCGGTTTCTGTATGCGAAGTTGCTAGGCCTGCTGGCTGA
GCTCCGGAGCATTAAAGAGCCTACGGGTACCAATCCAGCACATCCAGGCCTGTCTGCCATGATGCC
CTGCTCCAGGAGATCTGCAGC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA



[View online »](#)

Protein Sequence: >RC213905 representing NM_001077482
Red=Cloning site Green=Tags(s)

MASREDELRNCVCGDQATGYHFNALTCEGCKGFFRRTVSKSIGPTCPFAGSCEVSKTQRRHCPACRLQK
 CLDAGMRKDMILSAEALALRRAKQAQRRAQOTPVQLSKEQEELIRTLGHAHTRHMGTMFEQFVQFRPPAH
 LFIHHQPLPTLAPVLPVTHFADINTFMVLQVIKFTKDLPVFRSLPIEDQISLLKGAAVEICHIVLNTTF
 CLQTQNF LCGPLRYTIEDGARVSPTVGFQVEFLELLFHFHGTLRKLQLQEPEYVLLAAMALFSPAPYLTD
 RPGVTRQDEIDQLQEEMALTLQSYIKGQRRPRDRFLYAKLLGLLAELRSINEAYGYQIQHIQGLSAMMP
 LLQEICS

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Chromatograms: https://cdn.origene.com/chromatograms/mk6094_f07.zip

Restriction Sites: SgfI-MluI

Cloning Scheme:



ACCN: NM_001077482

ORF Size: 1071 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001077482.2](#)

RefSeq Size: 1408 bp

RefSeq ORF: 1074 bp

Locus ID: 9970

UniProt ID: [Q14994](#)


Cytogenetics: 1q23.3

Protein Families: Druggable Genome, Nuclear Hormone Receptor, Transcription Factors


MW: 40.3 kDa

Gene Summary: This gene encodes a member of the nuclear receptor superfamily, and is a key regulator of xenobiotic and endobiotic metabolism. The protein binds to DNA as a monomer or a heterodimer with the retinoid X receptor and regulates the transcription of target genes involved in drug metabolism and bilirubin clearance, such as cytochrome P450 family members. Unlike most nuclear receptors, this transcriptional regulator is constitutively active in the absence of ligand but is regulated by both agonists and inverse agonists. Ligand binding results in translocation of this protein to the nucleus, where it activates or represses target gene transcription. These ligands include bilirubin, a variety of foreign compounds, steroid hormones, and prescription drugs. In addition to drug metabolism, the CAR protein is also reported to regulate genes involved in glucose metabolism, lipid metabolism, cell proliferation, and circadian clock regulation. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2020]


Product images:



Circular map for RC213905



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY NR1I3 (Cat# RC213905, Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-NR1I3 (Cat# [TA805312]). Positive lysates [LY425868] (100ug) and [LC425868] (20ug) can be purchased separately from OriGene.



Western blot validation of overexpression lysate (Cat# [LY421437]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC213905 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).