

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for RC213918L3V

PAPOLG (NM_022894) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	PAPOLG (NM_022894) Human Tagged ORF Clone Lentiviral Particle
Symbol:	PAPOLG
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_022894
ORF Size:	2208 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC213918).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 022894.2</u>
RefSeq Size:	7386 bp
RefSeq ORF:	2211 bp
Locus ID:	64895
UniProt ID:	<u>Q9BWT3</u>
Cytogenetics:	2p16.1
Domains:	NTP_transf_2, PAP_RNA-bind, PAP_central
Protein Families:	Transcription Factors
Protein Pathways:	RNA degradation



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	PAPOLG (NM_022894) Human Tagged ORF Clone Lentiviral Particle – RC213918L3V
MW:	82.8 kDa
Gene Summary:	This gene encodes a member of the poly(A) polymerase family which catalyzes template- independent extension of the 3' end of a DNA/RNA strand. This enzyme shares 60% identity to the well characterized poly(A) polymerase II (PAPII) at the amino acid level. These two enzymes have similar organization of structural and functional domains. This enzyme is exclusively localized in the nucleus and exhibits both nonspecific and CPSF (cleavage and polyadenylation specificity factor)/AAUAAA-dependent polyadenylation activity. This gene is located on chromosome 2 in contrast to the PAPII gene, which is located on chromosome 14. [provided by RefSeq, Jul 2008]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US