

Product datasheet for RC213927L3

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

WISP3 (CCN6) (NM_003880) Human Tagged Lenti ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: WISP3 (CCN6) (NM_003880) Human Tagged Lenti ORF Clone

Tag: Myc-DDK

Symbol: CCN6

Synonyms: LIBC; PPAC; PPD; PPRD; WISP-3; WISP3

Mammalian Cell Puromycin

Selection:

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)


E. coli Selection: Chloramphenicol (34 ug/mL)

ORF Nucleotide The ORF insert of this clone is exactly the same as(RC213927).

Sgfl-Mlul

Sequence:

Restriction Sites: Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF.

ACCN: NM_003880

ORF Size: 1062 bp



WISP3 (CCN6) (NM_003880) Human Tagged Lenti ORF Clone - RC213927L3

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 003880.2</u>

 RefSeq Size:
 1307 bp

 RefSeq ORF:
 1065 bp

 Locus ID:
 8838

 UniProt ID:
 095389

Cytogenetics: 6q21

Protein Families: Druggable Genome, ES Cell Differentiation/IPS, Secreted Protein

MW: 39.1 kDa

Gene Summary: This gene encodes a member of the WNT1 inducible signaling pathway (WISP) protein

subfamily, which belongs to the connective tissue growth factor (CTGF) family. WNT1 is a member of a family of cysteine-rich, glycosylated signaling proteins that mediate diverse developmental processes. The CTGF family members are characterized by four conserved cysteine-rich domains: insulin-like growth factor-binding domain, von Willebrand factor type C module, thrombospondin domain and C-terminal cystine knot-like domain. This gene is overexpressed in colon tumors. It may be downstream in the WNT1 signaling pathway that is relevant to malignant transformation. Mutations of this gene are associated with progressive pseudorheumatoid dysplasia, an autosomal recessive skeletal disorder, indicating that the gene is essential for normal postnatal skeletal growth and cartilage homeostasis. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by

RefSeq, Jul 2008]