

Product datasheet for RC213959L4

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Macrophage Inflammatory Protein 1 beta (CCL4L2) (NM_001001435) Human Tagged Lenti ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: Macrophage Inflammatory Protein 1 beta (CCL4L2) (NM_001001435) Human Tagged Lenti ORF

Clone

Tag: mGFP

Symbol: Macrophage Inflammatory Protein 1 beta

Synonyms: AT744.2; CCL4L; LAG-1; LAG1; SCYA4L

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

E. coli Selection: Chloramphenicol (34 ug/mL)


ORF Nucleotide


The ORF insert of this clone is exactly the same as(RC213959).

Sequence:

Restriction Sites: Sgfl-Mlul

Cloning Scheme:





 $[\]ensuremath{^*}$ The last codon before the Stop codon of the ORF.

ACCN: NM 001001435





Macrophage Inflammatory Protein 1 beta (CCL4L2) (NM_001001435) Human Tagged Lenti ORF Clone - RC213959L4

ORF Size: 276 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001001435.2</u>, <u>NP 001001435.1</u>

RefSeq Size: 674 bp
RefSeq ORF: 278 bp
Locus ID: 9560
Cytogenetics: 17q12

Protein Families: Druggable Genome, Transmembrane

Protein Pathways: Chemokine signaling pathway, Cytokine-cytokine receptor interaction, Cytosolic DNA-sensing

pathway

MW: 10.1 kDa

Gene Summary: This gene is one of several cytokine genes that are clustered on the q-arm of chromosome 17.

Cytokines are a family of secreted proteins that function in inflammatory and


immunoregulatory processes. The protein encoded by this family member is similar to the chemokine (C-C motif) ligand 4 product, which inhibits HIV entry by binding to the cellular receptor CCR5. The copy number of this gene varies among individuals, where most

individuals have one to five copies. This gene copy contains a non-consensus splice acceptor site at the 3' terminal exon found in other highly similar gene copies, and it thus uses other alternative splice sites for the 3' terminal exon, resulting in multiple transcript variants.

[provided by RefSeq, Apr 2014]



Product images:



Circular map for RC213959L4