

Product datasheet for RC214065L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

p19 INK4d (CDKN2D) (NM 001800) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: p19 INK4d (CDKN2D) (NM_001800) Human Tagged ORF Clone Lentiviral Particle

Symbol: p19 INK4d

INK4D; p19; p19-INK4D Synonyms:

Mammalian Cell

Selection:

ACCN:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK NM 001800

ORF Size: 498 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC214065).

Sequence:

Cytogenetics:

The molecular sequence of this clone aligns with the gene accession number as a point of OTI Disclaimer:

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 001800.3, NP 001791.1

19p13.2

RefSeq Size: 1416 bp RefSeq ORF: 501 bp Locus ID: 1032 **UniProt ID:** P55273

Protein Families: Druggable Genome

Protein Pathways: Cell cycle





MW: 17.5 kDa

Gene Summary:

The protein encoded by this gene is a member of the INK4 family of cyclin-dependent kinase inhibitors. This protein has been shown to form a stable complex with CDK4 or CDK6, and prevent the activation of the CDK kinases, thus function as a cell growth regulator that controls cell cycle G1 progression. The abundance of the transcript of this gene was found to oscillate in a cell-cycle dependent manner with the lowest expression at mid G1 and a maximal expression during S phase. The negative regulation of the cell cycle involved in this protein was shown to participate in repressing neuronal proliferation, as well as spermatogenesis. Two alternatively spliced variants of this gene, which encode an identical protein, have been reported. [provided by RefSeq, Jul 2008]