

## Product datasheet for **RC214199L3V**

### **COPA (NM\_001098398) Human Tagged ORF Clone Lentiviral Particle**

#### **Product data:**

Product Type:	Lentiviral Particles
Product Name:	COPA (NM_001098398) Human Tagged ORF Clone Lentiviral Particle
Symbol:	COPA
Synonyms:	AILJK; alpha-COP; HEP-COP
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_001098398
ORF Size:	3699 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC214199).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<a href="#">NM_001098398.1</a> , <a href="#">NP_001091868.1</a>
RefSeq Size:	5666 bp
RefSeq ORF:	3702 bp
Locus ID:	1314
UniProt ID:	<a href="#">P53621</a>
Cytogenetics:	1q23.2
MW:	139.1 kDa



[View online »](#)

**Gene Summary:**

In eukaryotic cells, protein transport between the endoplasmic reticulum and Golgi compartments is mediated in part by non-clathrin-coated vesicular coat proteins (COPs). Seven coat proteins have been identified, and they represent subunits of a complex known as coatomer. The subunits are designated alpha-COP, beta-COP, beta-prime-COP, gamma-COP, delta-COP, epsilon-COP, and zeta-COP. The alpha-COP, encoded by COPA, shares high sequence similarity with RET1P, the alpha subunit of the coatomer complex in yeast. Also, the N-terminal 25 amino acids of alpha-COP encode the bioactive peptide, xenin, which stimulates exocrine pancreatic secretion and may act as a gastrointestinal hormone. Alternative splicing results in multiple splice forms encoding distinct isoforms. [provided by RefSeq, Jul 2008]