

## Product datasheet for RC214235L2V

### SEMA6D (NM\_153618) Human Tagged ORF Clone Lentiviral Particle

#### Product data:

Product Type:	Lentiviral Particles
Product Name:	SEMA6D (NM_153618) Human Tagged ORF Clone Lentiviral Particle
Symbol:	SEMA6D
Mammalian Cell Selection:	None
Vector:	pLenti-C-mGFP (PS100071)
Tag:	mGFP
ACCN:	NM_153618
ORF Size:	3219 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC214235).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<a href="#">NM_153618.1</a> , <a href="#">NP_705871.1</a>
RefSeq Size:	6109 bp
RefSeq ORF:	3222 bp
Locus ID:	80031
UniProt ID:	<a href="#">Q8NFY4</a>
Cytogenetics:	15q21.1
Protein Families:	Druggable Genome, Transmembrane
Protein Pathways:	Axon guidance
MW:	117.4 kDa


[View online »](#)

**Gene Summary:**

Semaphorins are a large family, including both secreted and membrane associated proteins, many of which have been implicated as inhibitors or chemorepellents in axon pathfinding, fasciculation and branching, and target selection. All semaphorins possess a semaphorin (Sema) domain and a PSI domain (found in plexins, semaphorins and integrins) in the N-terminal extracellular portion. Additional sequence motifs C-terminal to the semaphorin domain allow classification into distinct subfamilies. Results demonstrate that transmembrane semaphorins, like the secreted ones, can act as repulsive axon guidance cues. This gene encodes a class 6 vertebrate transmembrane semaphorin that demonstrates alternative splicing. Several transcript variants have been identified and expression of the distinct encoded isoforms is thought to be regulated in a tissue- and development-dependent manner. [provided by RefSeq, Nov 2010]