

Product datasheet for RC214315

CD89 (FCAR) (NM 133271) Human Tagged ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: CD89 (FCAR) (NM_133271) Human Tagged ORF Clone

Tag: Myc-DDK

Symbol: CD89

Synonyms: CD89; CTB-61M7.2; FcalphaRI

Mammalian Cell

Selection:

Neomycin

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)ORF Nucleotide>RC214315 ORF sequence

Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

AAGTGTCTGCAAG

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >RC214315 protein sequence

Red=Cloning site Green=Tags(s)

MDPKQTTLLCLVLCLGQRIQAQEGDFPMPFISAKSSPVIPLDGSVKIQCQAIREAYLTQLMIIKNSTYRE IGRRLKFWNETDPEFVIDHMDANKAGRYQCQYRIGHYRFRYSDTLELVVTDSIHQDYTTQNLIRMAVAGL

VLVALLAILVENWHSHTALNKEASADVAEPSWSQQMCQPGLTFARTPSVCK

TRTRPLEQKLISEEDLAANDILDYKDDDDKV



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Restriction Sites: Sgfl-Mlul

Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF

ACCN: NM_133271

ORF Size: 573 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 133271.2</u>, <u>NP 579805.1</u>

RefSeq Size: 1383 bp
RefSeq ORF: 576 bp
Locus ID: 2204
UniProt ID: P24071



Cytogenetics: 19q13.42

Protein Families: Transmembrane


MW: 21.7 kDa

Gene Summary: This gene is a member of the immunoglobulin gene superfamily and encodes a receptor for

the Fc region of IgA. The receptor is a transmembrane glycoprotein present on the surface of myeloid lineage cells such as neutrophils, monocytes, macrophages, and eosinophils, where it mediates immunologic responses to pathogens. It interacts with IgA-opsonized targets and triggers several immunologic defense processes, including phagocytosis, antibody-dependent cell-mediated cytotoxicity, and stimulation of the release of inflammatory mediators. Multiple alternatively spliced transcript variants encoding different isoforms have been described for

this gene. [provided by RefSeq, Jul 2008]

Product images:



Circular map for RC214315