

Product datasheet for RC214358L2V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

MEKK1 (MAP3K1) (NM_005921) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: MEKK1 (MAP3K1) (NM 005921) Human Tagged ORF Clone Lentiviral Particle

Symbol: MEKK1

Synonyms: MAPKKK1; MEKK; MEKK 1; MEKK1; SRXY6

Mammalian Cell

Selection:

None

Vector: pLenti-C-mGFP (PS100071)

Tag: mGFP

ACCN: NM_005921 **ORF Size:** 4536 bp

ORF Nucleotide

1330 66

Sequence:
OTI Disclaimer:

The ORF insert of this clone is exactly the same as(RC214358).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 005921.1

 RefSeq Size:
 7522 bp

 RefSeq ORF:
 4539 bp

 Locus ID:
 4214

 UniProt ID:
 Q13233

Cytogenetics: 5q11.2

Protein Families: Druggable Genome, Protein Kinase



MEKK1 (MAP3K1) (NM_005921) Human Tagged ORF Clone Lentiviral Particle - RC214358L2V

Protein Pathways: GnRH signaling pathway, MAPK signaling pathway, Neurotrophin signaling pathway, RIG-I-like

receptor signaling pathway, Ubiquitin mediated proteolysis

MW: 164.3 kDa

Gene Summary: The protein encoded by this gene is a serine/threonine kinase and is part of some signal

transduction cascades, including the ERK and JNK kinase pathways as well as the NF-kappa-B pathway. The encoded protein is activated by autophosphorylation and requires magnesium as a cofactor in phosphorylating other proteins. This protein has E3 ligase activity conferred by a plant homeodomain (PHD) in its N-terminus and phospho-kinase activity conferred by a

kinase domain in its C-terminus. [provided by RefSeq, Mar 2012]