

Product datasheet for RC214553L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

FZD3 (NM_017412) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: FZD3 (NM_017412) Human Tagged ORF Clone Lentiviral Particle

Symbol: FZD3
Synonyms: Fz-3

Mammalian Cell Puromycin

Selection:

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_017412 **ORF Size:** 1998 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC214553).

Sequence:

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 017412.2

RefSeq Size: 3933 bp
RefSeq ORF: 2001 bp
Locus ID: 7976
UniProt ID: Q9NPG1
Cytogenetics: 8p21.1

Domains: FRI, Frizzled

Protein Families: Druggable Genome, GPCR, Transmembrane





FZD3 (NM_017412) Human Tagged ORF Clone Lentiviral Particle - RC214553L4V

Protein Pathways: Basal cell carcinoma, Colorectal cancer, Melanogenesis, Pathways in cancer, Wnt signaling

pathway

MW: 76.1 kDa

Gene Summary: This gene is a member of the frizzled gene family. Members of this family encode seven-

transmembrane domain proteins that are receptors for the wingless type MMTV integration site family of signaling proteins. Most frizzled receptors are coupled to the beta-catenin canonical signaling pathway. The function of this protein is unknown, although it may play a role in mammalian hair follicle development. Alternative splicing results in multiple transcript variants. This gene is a susceptibility locus for schizophrenia. [provided by RefSeq, Dec 2010]