

## Product datasheet for **RC214570L1V**

### BAIAP2 (NM\_006340) Human Tagged ORF Clone Lentiviral Particle

#### Product data:

Product Type:	Lentiviral Particles
Product Name:	BAIAP2 (NM_006340) Human Tagged ORF Clone Lentiviral Particle
Symbol:	BAIAP2
Synonyms:	BAP2; FLAF3; IRSP53; WAML
Mammalian Cell Selection:	None
Vector:	pLenti-C-Myc-DDK (PS100064)
Tag:	Myc-DDK
ACCN:	NM_006340
ORF Size:	1560 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC214570).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<a href="#">NM_006340.1</a>
RefSeq Size:	2129 bp
RefSeq ORF:	1563 bp
Locus ID:	10458
UniProt ID:	<a href="#">Q9UQB8</a>
Cytogenetics:	17q25.3
Domains:	SH3
Protein Families:	Druggable Genome


[View online »](#)

**Protein Pathways:** Adherens junction, Regulation of actin cytoskeleton

**MW:** 57.3 kDa

**Gene Summary:** The protein encoded by this gene has been identified as a brain-specific angiogenesis inhibitor (BAI1)-binding protein. This adaptor protein links membrane bound G-proteins to cytoplasmic effector proteins. This protein functions as an insulin receptor tyrosine kinase substrate and suggests a role for insulin in the central nervous system. It also associates with a downstream effector of Rho small G proteins, which is associated with the formation of stress fibers and cytokinesis. This protein is involved in lamellipodia and filopodia formation in motile cells and may affect neuronal growth-cone guidance. This protein has also been identified as interacting with the dentatorubral-pallidoluysian atrophy gene, which is associated with an autosomal dominant neurodegenerative disease. Alternative splicing results in multiple transcript variants encoding distinct isoforms.[provided by RefSeq, Jan 2009]