

## Product datasheet for **RC214583L1V**

### Repulsive Guidance Molecule C (HFE2) (NM\_202004) Human Tagged ORF Clone Lentiviral Particle

#### Product data:

Product Type:	Lentiviral Particles
Product Name:	Repulsive Guidance Molecule C (HFE2) (NM_202004) Human Tagged ORF Clone Lentiviral Particle
Symbol:	Repulsive Guidance Molecule C
Synonyms:	HFE2; HFE2A; JH; RGM C
Mammalian Cell Selection:	None
Vector:	pLenti-C-Myc-DDK (PS100064)
Tag:	Myc-DDK
ACCN:	NM_202004
ORF Size:	600 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC214583).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<a href="#">NM_202004.2</a>
RefSeq Size:	1438 bp
RefSeq ORF:	603 bp
Locus ID:	148738
UniProt ID:	<a href="#">Q6ZVN8</a>
Cytogenetics:	1q21.1
Protein Families:	Transmembrane



[View online »](#)

**MW:** 21.3 kDa

**Gene Summary:** The product of this gene is involved in iron metabolism. It may be a component of the signaling pathway which activates hepcidin or it may act as a modulator of hepcidin expression. It could also represent the cellular receptor for hepcidin. Two uORFs in the 5' UTR negatively regulate the expression and activity of the encoded protein. Alternatively spliced transcript variants encoding different isoforms have been identified for this gene. Defects in this gene are the cause of hemochromatosis type 2A, also called juvenile hemochromatosis (JH). JH is an early-onset autosomal recessive disorder due to severe iron overload resulting in hypogonadotropic hypogonadism, hepatic fibrosis or cirrhosis and cardiomyopathy, occurring typically before age of 30. [provided by RefSeq, Oct 2015]