

## Product datasheet for RC214639L4V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## MALT1 (NM\_006785) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

**Product Name:** MALT1 (NM\_006785) Human Tagged ORF Clone Lentiviral Particle

Symbol: MALT1

Synonyms: IMD12; MLT; MLT1; PCASP1

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

**ACCN:** NM\_006785 **ORF Size:** 2472 bp

**ORF Nucleotide** 

- 1

Sequence:
OTI Disclaimer:

The ORF insert of this clone is exactly the same as(RC214639).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 006785.2</u>

RefSeq Size: 5029 bp
RefSeq ORF: 2475 bp
Locus ID: 10892
UniProt ID: Q9UDY8
Cytogenetics: 18q21.32

**Domains:** DEATH, ig, IGc2, IG

**Protein Families:** Druggable Genome, Protease





## MALT1 (NM\_006785) Human Tagged ORF Clone Lentiviral Particle - RC214639L4V

**Protein Pathways:** B cell receptor signaling pathway, T cell receptor signaling pathway

MW: 92.1 kDa

**Gene Summary:** This gene encodes a caspase-like protease that plays a role in BCL10-induced activation of

NF-kappaB. The protein is a component of the CARMA1-BCL10-MALT1 (CBM) signalosome that triggers NF-kappaB signaling and lymphoctye activation following antigen-receptor stimulation. Mutations in this gene result in immunodeficiency 12 (IMD12). This gene has been found to be recurrently rearranged in chromosomal translocations with other genes in mucosa-associated lymphoid tissue lymphomas, including a t(11;18)(q21;q21) translocation with the baculoviral IAP repeat-containing protein 3 (also known as apoptosis inhibitor 2) locus [BIRC3(API2)-MALT1], and a t(14;18)(q32;q21) translocation with the immunoglobulin heavy chain locus (IGH-MALT1). Alternatively spliced transcript variants have been described

for this gene. [provided by RefSeq, May 2018]