

### Product datasheet for RC214694L1

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Mu Opioid Receptor (OPRM1) (NM\_001008505) Human Tagged Lenti ORF Clone

**Product data:** 

**Product Type:** Expression Plasmids

Product Name: Mu Opioid Receptor (OPRM1) (NM\_001008505) Human Tagged Lenti ORF Clone

Tag: Myc-DDK

Symbol: Mu Opioid Receptor

Synonyms: LMOR; M-OR-1; MOP; MOR; MOR1; OPRM

Mammalian Cell None

Selection:

Vector:pLenti-C-Myc-DDK (PS100064)E. coli Selection:Chloramphenicol (34 ug/mL)

**ORF Nucleotide** The ORF insert of this clone is exactly the same as(RC214694).

Sequence:

**Restriction Sites:** Sgfl-Mlul

**Cloning Scheme:** 





<sup>\*</sup> The last codon before the Stop codon of the ORF

ACCN: NM\_001008505

ORF Size: 1338 bp



#### Mu Opioid Receptor (OPRM1) (NM\_001008505) Human Tagged Lenti ORF Clone - RC214694L1

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

**RefSeq:** <u>NM 001008505.1</u>, <u>NP 001008505.2</u>

6q25.2

 RefSeq Size:
 1652 bp

 RefSeq ORF:
 1341 bp

 Locus ID:
 4988

 UniProt ID:
 P35372

Cytogenetics:

**Protein Families:** Druggable Genome, GPCR, Transmembrane

**Protein Pathways:** Neuroactive ligand-receptor interaction

MW: 49.3 kDa

**Gene Summary:** This gene encodes one of at least three opioid receptors in humans; the mu opioid receptor

(MOR). The MOR is the principal target of endogenous opioid peptides and opioid analgesic agents such as beta-endorphin and enkephalins. The MOR also has an important role in dependence to other drugs of abuse, such as nicotine, cocaine, and alcohol via its modulation of the dopamine system. The NM\_001008503.2:c.118A>G allele has been associated with opioid and alcohol addiction and variations in pain sensitivity but evidence for it having a causal role is conflicting. Multiple transcript variants encoding different isoforms have been


found for this gene. Though the canonical MOR belongs to the superfamily of 7-

transmembrane-spanning G-protein-coupled receptors some isoforms of this gene have only


6 transmembrane domains. [provided by RefSeq, Oct 2013]



# **Product images:**



Circular map for RC214694L1



Double digestion of RC214694L1 using Sgfl and Mlul