

Product datasheet for RC214694L2V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Mu Opioid Receptor (OPRM1) (NM_001008505) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Mu Opioid Receptor (OPRM1) (NM 001008505) Human Tagged ORF Clone Lentiviral Particle

Symbol: Mu Opioid Receptor

Synonyms: LMOR; M-OR-1; MOP; MOR; MOR1; OPRM

Mammalian Cell

Selection:

None

Vector: pLenti-C-mGFP (PS100071)

Tag: mGFP

ACCN: NM_001008505

ORF Size: 1338 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC214694).

Sequence:

Cytogenetics:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 001008505.1, NP 001008505.2

 RefSeq Size:
 1652 bp

 RefSeq ORF:
 1341 bp

 Locus ID:
 4988

 UniProt ID:
 P35372

Protein Families: Druggable Genome, GPCR, Transmembrane

Protein Pathways: Neuroactive ligand-receptor interaction

6q25.2





Mu Opioid Receptor (OPRM1) (NM_001008505) Human Tagged ORF Clone Lentiviral Particle – RC214694L2V

MW: 49.3 kDa

Gene Summary:

This gene encodes one of at least three opioid receptors in humans; the mu opioid receptor (MOR). The MOR is the principal target of endogenous opioid peptides and opioid analgesic agents such as beta-endorphin and enkephalins. The MOR also has an important role in dependence to other drugs of abuse, such as nicotine, cocaine, and alcohol via its modulation of the dopamine system. The NM_001008503.2:c.118A>G allele has been associated with opioid and alcohol addiction and variations in pain sensitivity but evidence for it having a causal role is conflicting. Multiple transcript variants encoding different isoforms have been found for this gene. Though the canonical MOR belongs to the superfamily of 7-transmembrane-spanning G-protein-coupled receptors some isoforms of this gene have only 6 transmembrane domains. [provided by RefSeq, Oct 2013]