

Product datasheet for RC214833

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

CN: techsupport@origene.cn

OriGene Technologies, Inc.

TAF9 (NM 001015892) Human Tagged ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: TAF9 (NM 001015892) Human Tagged ORF Clone

Tag: Myc-DDK

Symbol: TAF9

Synonyms: MGC:5067; STAF31/32; TAF2G; TAFII-31; TAFII-32; TAFII31; TAFII32; TAFIID32

Mammalian Cell

Selection:

Neomycin

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)ORF Nucleotide>RC214833 ORF sequence

Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

 ${\tt TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC}$

GCCGCGATCGCC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATTACAAGGATGACGACGATAAGGTTTAA





Protein Sequence: >RC214833 protein sequence

Red=Cloning site Green=Tags(s)

MESGKTASPKSMPKDAQMMAQILKDMGITEYEPRVINQMLEFAFRYVTTILDDAKIYSSHAKKATVDADD VRLAIQCRADQSFTSPPPRDFLLDIARQRNQTPLPLIKPYSGPRLPPDRYCLTAPNYRLKSLQKKASTSA GRITVPRLSVGSVTSRPSTPTLGTPTPQTMSVSTKVGTPMSLTGQRFTVQMPTSQSPAVKASIPATSAVQ NVLINPSLIGSKNIFITTNMMSSQNTANESSNALKRKREDDDDDDDDDDDDDDDDDDNL


TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Chromatograms: https://cdn.origene.com/chromatograms/mk6770 a12.zip

Restriction Sites: Sgfl-Mlul

Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF

ACCN: NM_001015892

ORF Size: 792 bp

OTI Disclaimer: Due to the inherent nature of this plasmid, standard methods to replicate additional amounts

of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at customercom or by

calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.



TAF9 (NM_001015892) Human Tagged ORF Clone - RC214833

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001015892.1</u>, <u>NP 001015892.1</u>

 RefSeq Size:
 1482 bp

 RefSeq ORF:
 795 bp

 Locus ID:
 6880

 UniProt ID:
 Q16594

 Cytogenetics:
 5q13.2

Protein Families: Transcription Factors

Protein Pathways: Basal transcription factors

MW: 29 kDa

Gene Summary: Initiation of transcription by RNA polymerase II requires the activities of more than 70


polypeptides. The protein that coordinates these activities is transcription factor IID (TFIID), which binds to the core promoter to position the polymerase properly, serves as the scaffold for assembly of the remainder of the transcription complex, and acts as a channel for regulatory signals. TFIID is composed of the TATA-binding protein (TBP) and a group of evolutionarily conserved proteins known as TBP-associated factors or TAFs. TAFs may participate in basal transcription, serve as coactivators, function in promoter recognition or modify general transcription factors (GTFs) to facilitate complex assembly and transcription initiation. This gene encodes one of the smaller subunits of TFIID that binds to the basal transcription factor GTF2B as well as to several transcriptional activators such as p53 and VP16. In human, TAF9 and AK6 (GeneID: 102157402) are two distinct genes that share 5' exons. A similar but distinct gene (TAF9L) has been found on the X chromosome and a

pseudogene has been identified on chromosome 19. Alternative splicing results in multiple


transcript variants. [provided by RefSeq, Sep 2013]



Product images:



Circular map for RC214833



Western blot validation of overexpression lysate (Cat# [LY423137]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC214833 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).