

## Product datasheet for RC214837L2V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## PGPEP1 (NM\_017712) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

Product Name: PGPEP1 (NM\_017712) Human Tagged ORF Clone Lentiviral Particle

Symbol: PGPEP

**Synonyms:** PAP-I; Pcp; PGI; PGP-I; PGPI

**Mammalian Cell** 

Selection:

None

**Vector:** pLenti-C-mGFP (PS100071)

Tag: mGFP

**ACCN:** NM\_017712

ORF Size: 627 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC214837).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 017712.1</u>

 RefSeq Size:
 2239 bp

 RefSeq ORF:
 630 bp

 Locus ID:
 54858

 UniProt ID:
 Q9NXJ5

 Cytogenetics:
 19p13.11

**Protein Families:** Druggable Genome, Protease

MW: 23 kDa







## **Gene Summary:**

The gene encodes a cysteine protease and member of the peptidase C15 family of proteins. The encoded protein cleaves amino terminal pyroglutamate residues from protein substrates including thyrotropin-releasing hormone and other neuropeptides. Expression of this gene may be downregulated in colorectal cancer, while activity of the encoded protein may be negatively correlated with cancer progression in colorectal cancer patients. Activity of the encoded protease may also be altered in other disease states including in liver cirrhosis, which is associated with reduced protease activity, and in necrozoospermia, which is associated with elevated protease activity. [provided by RefSeq, Jul 2016]