

## Product datasheet for RC214873L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## CMA1 (NM\_001836) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** CMA1 (NM\_001836) Human Tagged ORF Clone Lentiviral Particle

Symbol: CMA1

**Synonyms:** chymase; CYH; MCT1

**Mammalian Cell** 

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

NM 001836

Tag: Myc-DDK

ORF Size: 741 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC214873).

Sequence:

ACCN:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 001836.1

 RefSeq Size:
 744 bp

 RefSeq ORF:
 744 bp

 Locus ID:
 1215

 UniProt ID:
 P23946

 Cytogenetics:
 14q12

**Protein Families:** Druggable Genome, Protease, Secreted Protein

**Protein Pathways:** Renin-angiotensin system







**MW:** 27.1 kDa

**Gene Summary:** 

This gene encodes a chymotryptic serine proteinase that belongs to the peptidase family S1. It is expressed in mast cells and is thought to function in the degradation of the extracellular matrix, the regulation of submucosal gland secretion, and the generation of vasoactive peptides. In the heart and blood vessels, this protein, rather than angiotensin converting enzyme, is largely responsible for converting angiotensin I to the vasoactive peptide angiotensin II. Alternative splicing results in multiple variants. [provided by RefSeq, Apr 2015]