

Product datasheet for RC214928L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

IGF1 Receptor (IGF1R) (NM 000875) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: IGF1 Receptor (IGF1R) (NM_000875) Human Tagged ORF Clone Lentiviral Particle

Symbol: IGF1 Receptor

Synonyms: CD221; IGFIR; IGFR; JTK13

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_000875 **ORF Size:** 4101 bp

ORF Nucleotide

Sequence:

The ORF insert of this clone is exactly the same as(RC214928).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 000875.2

 RefSeq Size:
 4989 bp

 RefSeq ORF:
 4104 bp

 Locus ID:
 3480

 UniProt ID:
 P08069

Cytogenetics: 15q26.3

Protein Families: Druggable Genome, Protein Kinase, Transmembrane





IGF1 Receptor (IGF1R) (NM_000875) Human Tagged ORF Clone Lentiviral Particle - RC214928L4V

Protein Pathways: Adherens junction, Colorectal cancer, Endocytosis, Focal adhesion, Glioma, Long-term

depression, Melanoma, Oocyte meiosis, Pathways in cancer, Progesterone-mediated oocyte

maturation, Prostate cancer

MW: 154.79 kDa

Gene Summary: This receptor binds insulin-like growth factor with a high affinity. It has tyrosine kinase

activity. The insulin-like growth factor I receptor plays a critical role in transformation events. Cleavage of the precursor generates alpha and beta subunits. It is highly overexpressed in most malignant tissues where it functions as an anti-apoptotic agent by enhancing cell survival. Alternatively spliced transcript variants encoding distinct isoforms have been found

for this gene. [provided by RefSeq, May 2014]