

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for RC214930L3V

## COPA (NM\_004371) Human Tagged ORF Clone Lentiviral Particle

## **Product data:**

Product Type:	Lentiviral Particles
Product Name:	COPA (NM_004371) Human Tagged ORF Clone Lentiviral Particle
Symbol:	СОРА
Synonyms:	AILJK; alpha-COP; HEP-COP
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_004371
ORF Size:	3672 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC214930).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 004371.3</u>
RefSeq Size:	5639 bp
RefSeq ORF:	3675 bp
Locus ID:	1314
UniProt ID:	<u>P53621</u>
Cytogenetics:	1q23.2
Domains:	WD40, Coatomer_WDAD
MW:	138.3 kDa



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



### COPA (NM\_004371) Human Tagged ORF Clone Lentiviral Particle – RC214930L3V

Gene Summary:In eukaryotic cells, protein transport between the endoplasmic reticulum and Golgi<br/>compartments is mediated in part by non-clathrin-coated vesicular coat proteins (COPs).<br/>Seven coat proteins have been identified, and they represent subunits of a complex known<br/>as coatomer. The subunits are designated alpha-COP, beta-COP, beta-prime-COP, gamma-<br/>COP, delta-COP, epsilon-COP, and zeta-COP. The alpha-COP, encoded by COPA, shares high<br/>sequence similarity with RET1P, the alpha subunit of the coatomer complex in yeast. Also, the<br/>N-terminal 25 amino acids of alpha-COP encode the bioactive peptide, xenin, which<br/>stimulates exocrine pancreatic secretion and may act as a gastrointestinal hormone.<br/>Alternative splicing results in multiple splice forms encoding distinct isoforms. [provided by<br/>RefSeq, Jul 2008]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US