

## Product datasheet for RC214967L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **ROR1 (NM 005012) Human Tagged ORF Clone Lentiviral Particle**

## **Product data:**

**Product Type: Lentiviral Particles** 

**Product Name:** ROR1 (NM 005012) Human Tagged ORF Clone Lentiviral Particle

Symbol:

dJ537F10.1; NTRKR1 Synonyms:

**Mammalian Cell** 

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK NM 005012 ACCN: **ORF Size:** 

**ORF Nucleotide** 

2811 bp

Sequence:

The ORF insert of this clone is exactly the same as(RC214967).

The molecular sequence of this clone aligns with the gene accession number as a point of OTI Disclaimer: reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

> clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 005012.1

RefSeq Size: 3358 bp RefSeq ORF: 2814 bp Locus ID: 4919 **UniProt ID:** Q01973 Cytogenetics: 1p31.3

**Domains:** KR, FRI, pkinase, TyrKc, S\_TKc, ig, IGc2, IG

**Protein Families:** Druggable Genome, Protein Kinase, Transmembrane





MW: 104.1 kDa

**Gene Summary:** 

This gene encodes a receptor tyrosine kinase-like orphan receptor that modulates neurite growth in the central nervous system. The encoded protein is a glycosylated type I membrane protein that belongs to the ROR subfamily of cell surface receptors. It is a pseudokinase that lacks catalytic activity and may interact with the non-canonical Wnt signalling pathway. This gene is highly expressed during early embryonic development but expressed at very low levels in adult tissues. Increased expression of this gene is associated with B-cell chronic lymphocytic leukaemia. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Jun 2012]