

## Product datasheet for RC215047L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## BMP8A (NM\_181809) Human Tagged ORF Clone Lentiviral Particle

## **Product data:**

Product Type: Lentiviral Particles

**Product Name:** BMP8A (NM\_181809) Human Tagged ORF Clone Lentiviral Particle

Symbol: BMP8A Synonyms: OP-2

Mammalian Cell Puromycin

Selection:

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

ACCN: NM\_181809

ORF Size: 1206 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC215047).

\_\_\_\_

Sequence:

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 181809.1

 RefSeq Size:
 1692 bp

 RefSeq ORF:
 1209 bp

 Locus ID:
 353500

 UniProt ID:
 Q7Z5Y6

 Cytogenetics:
 1p34.3

**Protein Families:** Secreted Protein

**Protein Pathways:** Hedgehog signaling pathway, TGF-beta signaling pathway





ORIGENE

**MW:** 44.6 kDa

**Gene Summary:** 

This gene encodes a secreted ligand of the TGF-beta (transforming growth factor-beta) superfamily of proteins. Ligands of this family bind various TGF-beta receptors leading to recruitment and activation of SMAD family transcription factors that regulate gene expression. The encoded preproprotein is proteolytically processed to generate each subunit of the disulfide-linked homodimer. This protein may play a role in development of the reproductive system. This gene may have arose from a gene duplication event and its gene duplicate is also present on chromosome 1. [provided by RefSeq, Jul 2016]