

## Product datasheet for RC215158L1V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **EGLN1 (NM 022051) Human Tagged ORF Clone Lentiviral Particle**

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** EGLN1 (NM\_022051) Human Tagged ORF Clone Lentiviral Particle

Symbol:

C1orf12; ECYT3; HALAH; HIF-PH2; HIFPH2; HPH-2; HPH2; PHD2; SM20; ZMYND6 Synonyms:

**Mammalian Cell** 

Selection:

None

Vector: pLenti-C-Myc-DDK (PS100064)

Myc-DDK Tag: NM 022051 ACCN: **ORF Size:** 1278 bp

**ORF Nucleotide** 

Sequence:

The ORF insert of this clone is exactly the same as(RC215158).

The molecular sequence of this clone aligns with the gene accession number as a point of OTI Disclaimer: reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 022051.1

RefSeq Size: 7102 bp RefSeq ORF: 1281 bp Locus ID: 54583 **UniProt ID:** Q9GZT9 Cytogenetics: 1q42.2

Domains: zf-MYND, 2OG-FeII\_Oxy, P4Hc

**Protein Pathways:** Pathways in cancer, Renal cell carcinoma





## EGLN1 (NM\_022051) Human Tagged ORF Clone Lentiviral Particle - RC215158L1V

**MW:** 46.5 kDa

**Gene Summary:** The protein encoded by this gene catalyzes the post-translational formation of 4-

hydroxyproline in hypoxia-inducible factor (HIF) alpha proteins. HIF is a transcriptional complex that plays a central role in mammalian oxygen homeostasis. This protein functions as a cellular oxygen sensor, and under normal oxygen concentration, modification by prolyl hydroxylation is a key regulatory event that targets HIF subunits for proteasomal destruction via the von Hippel-Lindau ubiquitylation complex. Mutations in this gene are associated with

erythrocytosis familial type 3 (ECYT3). [provided by RefSeq, Nov 2009]