

#### OriGene Technologies, Inc.


9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for RC215219L2

### Somatostatin Receptor 3 (SSTR3) (NM\_001051) Human Tagged Lenti ORF Clone

#### **Product data:**

| Product Type:                | Expression Plasmids  |
|------------------------------|--|
| Product Name:                | Somatostatin Receptor 3 (SSTR3) (NM_001051) Human Tagged Lenti ORF Clone                   |
| Tag:                         | mGFP   |
| Symbol:                      | Somatostatin Receptor 3  |
| Synonyms:                    | SS-3-R; SS3-R; SS3R; SSR-28  |
| Mammalian Cell<br>Selection: | None   |
| Vector:                      | pLenti-C-mGFP (PS100071)   |
| E. coli Selection:           | Chloramphenicol (34 ug/mL)   |
| ORF Nucleotide<br>Sequence:  | The ORF insert of this clone is exactly the same as(RC215219).                             |
| <b>Restriction Sites:</b>    | Sgfl-Mlul  |
| Cloning Scheme:              |  |
|                              | Cloning sites used for ORF Shuttling:  |
|                              | Sgf I         ORF         Mlu I            GCG ATC GC C         ATG // NNN         ACG CGT |



ACCN: ORF Size: NM\_001051 1254 bp




View online »

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

|                        | ostatin Receptor 3 (SSTR3) (NM_001051) Human Tagged Lenti ORF Clone – RC215219L2   |
|------------------------|--|
| OTI Disclaimer:        | The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>  |
| OTI Annotation:        | This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.   |
| Components:            | The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).   |
| Reconstitution Method: | <ol> <li>Centrifuge at 5,000xg for 5min.</li> <li>Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li> <li>Close the tube and incubate for 10 minutes at room temperature.</li> <li>Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li> <li>Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li> </ol>   |
| RefSeq:                | <u>NM 001051.2</u>   |
| RefSeq Size:           | 2123 bp  |
| RefSeq ORF:            | 1257 bp  |
| Locus ID:              | 6753   |
| UniProt ID:            | <u>P32745</u>  |
| Cytogenetics:          | 22q13.1  |
| Protein Families:      | Druggable Genome, GPCR, Transmembrane  |
| Protein Pathways:      | Neuroactive ligand-receptor interaction  |
| MW:                    | 45.7 kDa   |
| Gene Summary:          | This gene encodes a member of the somatostatin receptor protein family. Somatostatins are peptide hormones that regulate diverse cellular functions such as neurotransmission, cell proliferation, and endocrine signaling as well as inhibiting the release of many hormones and other secretory proteins. Somatostatin has two active forms of 14 and 28 amino acids. The biological effects of somatostatins are mediated by a family of G-protein coupled somatostatin receptors that are expressed in a tissue-specific manner. Somatostatin receptors form homodimers and heterodimers with other members of the superfamily as well as with other G-protein coupled receptors and receptor tyrosine kinases. This protein is functionally coupled to adenylyl cyclase. Alternate splicing results in multiple transcript variants. [provided by RefSeq, Jul 2013] |

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

## **Product images:**



Circular map for RC215219L2

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US