

## Product datasheet for **RC215275**

### SRPK2 (NM\_182692) Human Tagged ORF Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	SRPK2 (NM_182692) Human Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	SRPK2
Synonyms:	SFRSK2
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)

**ORF Nucleotide Sequence:**

>RC215275 representing NM\_182692  
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTGACTGGATCCGGTACCGAGGAGATCTGCC  
 GCC**CGGATCGCC**

ATGAGCTCCCGAAAGTGTGGCCATTCAGGCCGAAAGCGGAGGCCGAAAAGAGAGAAACATCCGAAAA  
 AGCCGGAGCCTCAACAGAAAGCTCCTTTAGTTCCTCCTCCACCGCCACCACCACCACCACCACC  
 TTTGCCAGACCCACACCCCGGAGCCAGAGGAGGAGATCCTGGGATCAGATGATGAGGACCAAGAGGAC  
 CCTGCGGACTACTGCAAAGGTGGATATCATCCAGTGAAAATTGGAGACCTCTTCAATGGCCGGTATCATG  
 TTATTAGAAAGCTTGGATGGGGCACTTCTCTACTGTCTGGTGTGCTGGGATATGCAGGGGAAAAGATT  
 TGTTGCAATGAAAGTTGTAAGTGGCCAGCATTATACGGAGACAGCCTTGGATGAAATAAAATTGCTC  
 AAATGTGTTGAGAAAGTATCCAGTGACCCAAACAAAGACATGGTGGTCCAGCTCATTGACGACTTCA  
 AGATTCAGGCATGAATGGGATACATGTCTGCATGGTCTTGAAGTACTTGGCCACCATCTCCTCAAGTG  
 GATCATCAAATCCAATCAAGGCCTCCAGTACGTTGTGTAAGAGTATCATTGACAGGTCCTTCAA  
 GGGTTAGATTACTTACACAGTAAGTGAAGATCATTCACTGACATAAAGCCGGAAAATATCTTGATGT  
 GTGTGGATGATGCATATGTGAGAAGAATGGCAGCTGAGGCCACTGAGTGGCAGAAAAGCAGGTGCTCTCC  
 TCCTTCAGGGTCTGCAGTGAGTACGGCTCCACAGCAGAAACCTATAGGAAAAATATCTAAAAACAAAAG  
 AAAAACTGAAAAAGAAACAGAAAGGCCAGGCTGAGTTATTGGAGAAGCGCTGCAGGAGATAGAAGAAT  
 TGGAGCGAGAAGCTGAAAGGAAAATAATAGAAGAAAACATCACCTCAGCTGCACCTTCCAATGACCAGGA  
 TGGCGAATACTGCCAGAGGTGAACTAAAAACAACAGGATTAGAGGAGGCGGCTGAGGCAGAGACTGCA  
 AAGGACAATGGTGAAGCTGAGGACCAGGAAGAGAAAGAAGATGCTGAGAAAGAAAACATTTGAAAAGATG  
 AAGATGATGTAGATCAGGAACCTGCGAACATAGACCCTACGTGGATAGAATCACCTAAAACCAATGGCCA  
 TATTGAGAAATGGCCATTCTCACTGGAGCAGCAACTGGACGATGAAGATGATGATGAAGAAGACTGCCCA  
 AATCCTGAGGAATATAATCTTGATGAGCCAAATGCAGAAAGTATTACACATATAGCAGCTCCTATGAAC  
 AATTCAATGGTGAATTGCCAAATGGACGACATAAAATTCCCGAGTACAGTTCACAGATTTCCACCTC  
 GTTGTCTCTGGATCCTTAGAACCTGTGGCCTGCGGCTCTGTGCTTTCTGAGGGATCACCCTTACTGAG  
 CAAGAGGAGAGCAGTCCATCCCATGACAGAAGCAGAACGGTTTCAGCCTCCAGTACTGGGGATTTGCCAA  
 AAGCAAAAACCCGGCAGCTGACTTGTGGTGAATCCCCTGGATCCGCGGAATGCAGATAAAATTAGAGT  
 AAAAATTGCTGACCTGGGAAATGCTTGTGGGTGCATAAACACTTCACGGAAGACATCCAGACGCGTCAG  
 TACCGCTCCATAGAGTTTTAATAGGAGCGGGTACAGCACCCCTGCGGACATCTGGAGCACGGCGTGTA  
 TGGCATTGAGCTGGCAACGGGAGATTATTTGTTGAACCACATTCTGGGGAAGACTATTCCAGAGACGA  
 AGACCACATAGCCACATCATAGAGCTGTAGGCAGTATTCCAAGGCACTTTGCTCTATCTGAAAAATAT  
 TCTCGGAATTCTTCAATCGCAGAGGAGAACTGCGACACATCACCAAGCTGAAGCCCTGGAGCCTTTTG  
 ATGTAATTGTGAAAAAGTATGGCTGGCCCATGAAGATGCTGCACAGTTTACAGATTTCTGATCCCGAT  
 GTTAGAAATGTTCCAGAAAACGAGCCTCAGCTGGCGAATGCCTTCGGCATCCTTGTTGAATTCT

AG**CGGACCG**ACGCGTACGCGGCCGCTCGAGCAGAAAACCTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCC  
 TGGATTACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >RC215275 representing NM\_182692  
 Red=Cloning site Green=Tags(s)


MSSRKVLAIQARKRRPKREKHPKKPEPQQKAPLVPPPPPPPPPPPLDPTPEPEEEEILGSDDEEQED  
 PADYCKGGYHPVKIGDLFNGRYHVIRKLGWGHFSTVWLCWDMQGRFVAMKVVKSAQHYTETALDEIKLL  
 KCVRESDPKDMVQVLDLDFKISGMNGIHVCMVFEVLGHLLKWIISKSNYQGLPVRCVKSIIRQVQLQ  
 GLDYLHSHKCKIIHTDIKPENILMCVDDAYVRRMAAEATEWQKAGAPPPSGSAVSTAPQPKPIGKISKNNK  
 KKLKQKQRQAELLEKRLEIEELEREAEERKIIIEENITSAAPSNQDGEYCPEVKLKTGLLEAAEAETA  
 KDNGEAEDQEEKEDAENIEKDEDDVDQELANIDPTWIESPKTNGHIENGPFSLEQQLDDEDDDEEDCP  
 NPPEYNLDEPNAESDYTYSSSYEQFNGELPNGRHKIPESQFPEFSTSLFSGSLEPVACGSVLSEGSPLTE  
 QEESSPHDRSRTVSASSTGDLPAKTRAADLLVNPLDPRNADKIRVKIADLGNACWVHKHFTEDIQTRQ  
 YRSIEVLIGAGYSTPADIWSTACMAFELATGDYLFEPHSGEDYSRDEDHIAHIELLGSI PRHFALSGKY  
 SREFFNRRGELRHITKLPWSLFDVLEKYGWPHEDAAQFDLIPMLEMVPEKRASAGECLRHPWLNLS

SGP TRRRLEQKLI SEEDLAANDILDYKDDDDKV

Chromatograms: [https://cdn.origene.com/chromatograms/mk8073\\_d10.zip](https://cdn.origene.com/chromatograms/mk8073_d10.zip)

Restriction Sites: SgfI-RsrII

Cloning Scheme:



ACCN: NM\_182692


ORF Size: 2097 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)


OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<a href="#">NM_182692.3</a>
<b>RefSeq Size:</b>	3671 bp
<b>RefSeq ORF:</b>	2100 bp
<b>Locus ID:</b>	6733
<b>UniProt ID:</b>	<a href="#">P78362</a>
<b>Cytogenetics:</b>	7q22.3
<b>Protein Families:</b>	Druggable Genome, Protein Kinase
<b>MW:</b>	78.8 kDa
<b>Gene Summary:</b>	Serine/arginine-rich protein-specific kinase which specifically phosphorylates its substrates at serine residues located in regions rich in arginine/serine dipeptides, known as RS domains and is involved in the phosphorylation of SR splicing factors and the regulation of splicing. Promotes neuronal apoptosis by up-regulating cyclin-D1 (CCND1) expression. This is done by the phosphorylation of SRSF2, leading to the suppression of p53/TP53 phosphorylation thereby relieving the repressive effect of p53/TP53 on cyclin-D1 (CCND1) expression. Phosphorylates ACIN1, and redistributes it from the nuclear speckles to the nucleoplasm, resulting in cyclin A1 but not cyclin A2 up-regulation. Plays an essential role in spliceosomal B complex formation via the phosphorylation of DDX23/PRP28. Can mediate hepatitis B virus (HBV) core protein phosphorylation. Plays a negative role in the regulation of HBV replication through a mechanism not involving the phosphorylation of the core protein but by reducing the packaging efficiency of the pregenomic RNA (pgRNA) without affecting the formation of the viral core particles.[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for RC215275



Western blot validation of overexpression lysate (Cat# [LY405411]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC215275 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).