

## Product datasheet for RC215286L1V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Lipin 3 (LPIN3) (NM\_022896) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

Product Name: Lipin 3 (LPIN3) (NM\_022896) Human Tagged ORF Clone Lentiviral Particle

Symbol: Lipin 3

**Synonyms:** dJ620E11.2; LIPN3L; SMP2

**Mammalian Cell** 

Selection:

None

**Vector:** pLenti-C-Myc-DDK (PS100064)

 Tag:
 Myc-DDK

 ACCN:
 NM\_022896

ORF Size: 2553 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC215286).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeq:** <u>NM 022896.1</u>

 RefSeq Size:
 4464 bp

 RefSeq ORF:
 2556 bp

 Locus ID:
 64900

 Cytogenetics:
 20q12

 MW:
 93.4 kDa





## **Gene Summary:**

The protein encoded by this gene is a member of the lipin family of proteins, and all family members share strong homology in their C-terminal region. This protein is thought to form hetero-oligomers with other lipin family members, while one family member, lipin 1, can also form homo-oligomers. This protein contains conserved motifs for phosphatidate phosphatase 1 (PAP1) activity as well as a domain that interacts with a transcriptional coactivator. Lipin complexes act in the cytoplasm to catalyze the dephosphorylation of phosphatidic acid to produce diacylglycerol, which is the precursor of both triglycerides and phospholipids. Lipin complexes are also thought to regulate gene expression as transcriptional co-activators in the nucleus. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Sep 2014]