

## Product datasheet for RC215290L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## PATZ1 (NM\_032050) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

**Product Name:** PATZ1 (NM\_032050) Human Tagged ORF Clone Lentiviral Particle

Symbol: PATZ1

Synonyms: dJ400N23; MAZR; PATZ; RIAZ; ZBTB19; ZNF278; ZSG

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

 Tag:
 Myc-DDK

 ACCN:
 NM\_032050

 ORF Size:
 1923 bp

**ORF Nucleotide** 

OTI Disclaimer:

Sequence:

The ORF insert of this clone is exactly the same as(RC215290).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 032050.1

 RefSeq Size:
 3674 bp

 RefSeq ORF:
 1926 bp

 Locus ID:
 23598

 UniProt ID:
 Q9HBE1

 Cytogenetics:
 22q12.2

**Domains:** BTB, AT\_hook, zf-C2H2

**Protein Families:** Transcription Factors





ORIGENE

**MW:** 68.9 kDa

**Gene Summary:** 

The protein encoded by this gene contains an A-T hook DNA binding motif which usually binds to other DNA binding structures to play an important role in chromatin modeling and transcription regulation. Its Poz domain is thought to function as a site for protein-protein interaction and is required for transcriptional repression, and the zinc-fingers comprise the DNA binding domain. Since the encoded protein has typical features of a transcription factor, it is postulated to be a repressor of gene expression. In small round cell sarcoma, this gene is fused to EWS by a small inversion of 22q, then the hybrid is thought to be translocated (t(1;22)(p36.1;q12). The rearrangement of chromosome 22 involves intron 8 of EWS and exon 1 of this gene creating a chimeric sequence containing the transactivation domain of EWS fused to zinc finger domain of this protein. This is a distinct example of an intrachromosomal rearrangement of chromosome 22. Four alternatively spliced transcript variants are described for this gene. [provided by RefSeq, Jul 2008]