

Product datasheet for RC215308

HGF (NM_001010933) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	HGF (NM_001010933) Human Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	HGF
Synonyms:	DFNB39; F-TCF; HGFB; HPTA; SF
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
ORF Nucleotide Sequence:	>RC215308 representing NM_001010933 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGGATCGCC**

ATGTGGGTGACCAAACCTCTGCCAGCCCTGCTGCTGCAGCATGTCCTCCTGCATCTCCTCCTGCTCCCA
TCGCCATCCCCTATGCAGAGGGACAAAGGAAAAGAAGAAATACAATTCATGAATCAAAAAATCAGCAA
GACTACCCTAATCAAAATAGATCCAGCACTGAAGATAAAAACCAAAAAAGTGAATACTGCAGACCAATGT
GCTAATAGATGTACTAGGAATAAAGGACTTCCATTCCTTCAAGGCTTTTGTGTTTGTAAAGCAAGAA
AACAAATGCCTCTGGTTCCCTTCAATAGCATGTCAAGTGGAGTAAAAAAGAATTTGGCCATGAATTTGA
CCTCTATGAAAACAAAGACTACATTAGAACTGCATCATTGGTAAAGGACGCAGCTACAAGGAACAGTA
TCTATCACTAAGAGTGGCATCAAATGTCAGCCCTGGAGTTCATGATACACACGAACACAGCTATCGGG
GTAAAGACCTACAGGAAACTACTGTGCAATCCTCGAGGGGAAGAAGGGGGACCCTGGTGTTCACAAG
CAATCCAGAGGTACGCTACGAAGTCTGTGACATTCCTCAGTGTTCAGAAGTTGAATGCATGACCTGCAAT
GGGAGAGTTATCGAGGTCTCATGGATCATACAGAATCAGGCAAGATTTGTCAGCGCTGGGATCATCAGA
CACCACACCGGCACAAATCTTGCCTGAAAGATATCCCGACAAGGGCTTTGATGATAATTATTGCCGCAA
TCCCGATGGCCAGCCGAGGCCATGGTGTATACTCTTGACCCTCACACCCGCTGGGAGTACTGTGCAATT
AAAACATGCGAGACA

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA



[View online »](#)

Protein Sequence: >RC215308 representing NM_001010933
Red=Cloning site Green=Tags(s)

MWVTKLLPALLLQHVLLHLLLLPIAIPYAEGQRKRRTIHEFKKSAKTTLIKIDPALKIKTKKVNTADQC
 ANRCTRNGLPFTCKAFVFDKARKQCLWFPFNSMSSGVKKEFGHEFDLYENKDYIRNCIIGKGRSYKGTV
 SITKSGIKQPWSSMIPHEHSYRGLDQENYCRNPRGEEGGPWCFSTSNPEVRYEVCIPQCSEVECMTCN
 GESYRGLMDHTESGKICQRWDHQTPHRHKFLPERYPDKGFDDNYCRNPDGQPRPWCYTLDPHTRWEYCAI
 KT CET

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Chromatograms: https://cdn.origene.com/chromatograms/mk8049_d04.zip

Restriction Sites: SgfI-MluI

Cloning Scheme:



ACCN: NM_001010933

ORF Size: 855 bp


OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.


Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_001010933.3
RefSeq Size:	1292 bp
RefSeq ORF:	858 bp
Locus ID:	3082
UniProt ID:	P14210
Cytogenetics:	7q21.11
Protein Families:	Adult stem cells, Druggable Genome, ES Cell Differentiation/IPS, Protease, Transmembrane
Protein Pathways:	Cytokine-cytokine receptor interaction, Focal adhesion, Melanoma, Pathways in cancer, Renal cell carcinoma
MW:	33.23 kDa
Gene Summary:	<p>This gene encodes a protein that binds to the hepatocyte growth factor receptor to regulate cell growth, cell motility and morphogenesis in numerous cell and tissue types. Alternative splicing results in multiple transcript variants, at least one of which encodes a preproprotein that is proteolytically processed to generate alpha and beta chains, which form the mature heterodimer. This protein is secreted by mesenchymal cells and acts as a multi-functional cytokine on cells of mainly epithelial origin. This protein also plays a role in angiogenesis, tumorigenesis, and tissue regeneration. Although the encoded protein is a member of the peptidase S1 family of serine proteases, it lacks peptidase activity. Mutations in this gene are associated with nonsyndromic hearing loss. [provided by RefSeq, Nov 2015]</p>

Product images:



Circular map for RC215308



Western blot validation of overexpression lysate (Cat# [LY423238]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC215308 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).