

## Product datasheet for RC215308L2V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## HGF (NM\_001010933) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** HGF (NM\_001010933) Human Tagged ORF Clone Lentiviral Particle

Symbol: HGF

Synonyms: DFNB39; F-TCF; HGFB; HPTA; SF

Mammalian Cell

Selection:

None

**Vector:** pLenti-C-mGFP (PS100071)

Tag: mGFP

**ACCN:** NM\_001010933

ORF Size: 855 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC215308).

Sequence:

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 001010933.1

 RefSeq Size:
 1292 bp

 RefSeq ORF:
 858 bp

 Locus ID:
 3082

 UniProt ID:
 P14210

 Cytogenetics:
 7q21.11

**Protein Families:** Adult stem cells, Druggable Genome, ES Cell Differentiation/IPS, Protease, Transmembrane





## HGF (NM\_001010933) Human Tagged ORF Clone Lentiviral Particle - RC215308L2V

**Protein Pathways:** Cytokine-cytokine receptor interaction, Focal adhesion, Melanoma, Pathways in cancer, Renal

cell carcinoma

MW: 33.23 kDa

**Gene Summary:** This gene encodes a protein that binds to the hepatocyte growth factor receptor to regulate

cell growth, cell motility and morphogenesis in numerous cell and tissue types. Alternative splicing results in multiple transcript variants, at least one of which encodes a preproprotein that is proteolytically processed to generate alpha and beta chains, which form the mature heterodimer. This protein is secreted by mesenchymal cells and acts as a multi-functional cytokine on cells of mainly epithelial origin. This protein also plays a role in angiogenesis, tumorogenesis, and tissue regeneration. Although the encoded protein is a member of the peptidase S1 family of serine proteases, it lacks peptidase activity. Mutations in this gene are

associated with nonsyndromic hearing loss. [provided by RefSeq, Nov 2015]